### The interatrial blocks versus atrial enlargements

### Dr. Andrés R. Pérez Riera

### Overview

IAB exists as an anatomical-electrical entity, which should be considered a true block. IAB presents with different degrees as other blocks in the conduction system. It shows a correlation with the left atrium size, however, it can be seen in patients with normal atrial size too. IAB is strongly associated with AF, and PACs and it could be considered a predictor of cardioembolic stroke. IAB is an expression of atrial electrical remodeling and dysfunction. IAB can be transient and in certain clinical circumstances, may be reversible. The contribution of endocardial mapping has increased our knowledge of the anatomy and pathophysiology of IAB.

The IAB classification should include first, second and advanced or third degree IAB. The P wave morphology should always be taking into consideration when diagnosing this condition. Finally, without the initial description of IAB made by Dr. Bayés de Luna, it would be impossible to understand IAB as an anatomical and electrical substrate for atrial arrhythmias. This represents a major contribution to the knowledge of electrocardiography and electrophysiology, and makes commendable that this arrhythmic syndrome should be called «Bayés' syndrome»(Conde 2014; 2015)

## I) First-degree (Partial IAB)

The electrical impulse is conducted from the right atrium (RA) to the left atrium (LA) trough Bachmann's bundle, but conduction is delayed. The ECG shows a P wave of  $\geq$  110ms in several leads with a variable negative wave in V<sub>1</sub>. The P-wave morphology is similar to that in left atrial enlargement, but usually a negative P wave in V<sub>1</sub> is less evident.



### Prevalence

High. Several studies have reported that the prevalence of IAB is more than 40% in hospital inpatients. Despite this, IAB remains largely underdiagnosed and commonly ignored. Advanced form is much less common that partial form (Kitkungvan 2009). Spodic consider this ECG abnormality is present in pandemic proportions especially at ages 60 and over and in unselected hospital patients. Because of its pathologic implications it requires widespread attention as a "pandemy". (Spodic 2009)

#### Associated conditions

- 1. Coronary Artery Disease
- 2. Hypertension
- 3. Diabetes Mellitus
- 4. Atrial fibrillation (strong associations)
- 5. Potential risk for embolism
- 6. Left Atrial Enlargement(LAE).
- 7. Left atrial electromechanical dysfunction.

### **Electrocardiographic characterization**

- P-wave duration ≥ 120ms(Ariyarajah 2006). P-wave duration is generally accepted as the most reliable non-invasive marker of atrial conduction and its prolongation is stronger associated with AF. However, patients with paroxysmal AF without structural heart disease may not have P-wave prolongation thus suggesting that the global conduction slowing is not an obligatory requirement for development of AF. (Platonov 2008)
- 2) P terminal force (Ptf) plus-minus P wave (biphasic configuration) in lead V1≥ the area of one small square the final minus portion indicates left atrial abnormality, particularly LAE, which is a strong correlate of IAB.
- 3) Prolonged in
- 4) trinsecoid P-wave deflection (from the apex to nadir) of the biphasic P wave in lead  $V1 \ge 40$ ms.
- 5) Often bifid ("notched") P waves
- 6) "Dome-and-spike" P-waves. 4 and 5 predominantly on leads II and from V<sub>3</sub> to V<sub>6</sub>.
  (Ariyarajah2006)

P-wave morphology and duration reveals several aspects of the atria: Proper function, fibrosis, dyssynchrony, final diastolic end LV pressure and activation paths can be inferred from the surface P-wave analysis. ECG can help differentiating atrial enlargements from conduction defects including intra- and IAB. (Baranchuk 20015)

Representation of normal P wave in V1



partial IAB is not associated with LAE.

For some authors the name "atrial abnormalities" encompass the concept of atrial enlargement and IABs (**Bayés de Luna 2013**)( Lee 2007). Both concepts, emphasizing, as it happens with ventricular hypertrophy and ventricular blocks, that often the ECG/VCG pattern of atrial enlargement especially of left atrial enlargement (LAE) is explained by the coexistence of IAB. However there are clear evidences that the pattern of IAB may exist without the association of atrial enlargement/hypertrophy. Therefore IABs and atrial enlargement are separate entities that are often associated with each other. Therefore we do not consider it appropriate to use the "umbrella" term atrial abnormalities to include both concepts without distinguishing between them (Lee 2007), as some authors do (**Bayés de Luna 2013**: ) (**Tsao 2008**).

#### Treatment

Pharmacological approach: Angiotensin-converting enzyme inhibitors and Angiotensin receptor blockers

Pacing

#### Advanced interatrial block or third degree interatrial block

Electrical impulse is blocked in Bachmann's bundle, but retrograde left atrial activation usually occurs.



Bayés de Luna A et al (**Bayés de Luna 1988**) studied 16 patients with ECG evidence of advanced interatrial block and retrograde activation of the left atrium (LA): P duration  $\geq$  120ms, and plus-minus biphasic (+/-) P waves in inferior leads II, III, and VF.

Eight patients had valvar heart disease, four had dilated cardiomyopathy and four had other forms of heart disease.

Patients with valvar heart disease and cardiomyopathy were compared with a control group of 22 patients with similar clinical and echocardiographic characteristics, but without this type of interatrial block.

Patients with advanced interatrial block and retrograde activation of the LA had a much higher incidence of paroxysmal supraventricular tachyarrhythmia's (93.7%) during followup than did the control group. Eleven of 16 patients (68.7%) with advanced interatrial block and retrograde activation of LA had atrial flutter (atypical in seven cases, typical in two cases, and with two or more morphologies in two cases). Six patients from the control group (27.7%) had sustained atrial tachyarrhythmias (five atrial fibrillation and one typical atrial flutter). The atrial tachyarrhythmias were due more to advanced interatrial block and retrograde activation of LA and frequent PACs than to LAE, because the control group with a LA of the same size, but without advanced interatrial block and retrograde activation of LA and frequent PACs, had a much lower incidence of paroxysmal tachycardia.

Bayés de Luna et al( **Bayés de Luna 1989**) demonstrated the value of preventive antiarrhythmic treatment in patients with advanced interatrial block

Diagnosis criteria of advanced interatrial block and retrograde activation of the left atrium(Bayes de Luna 1977;1988)

- 1. Biphasic P waves, in inferior leads.
- 2.  $P duration \ge 120 ms$
- 3. Angle between the first portion (RA) and end portion (LA) >90°
- 4. Orthogonal Y lead plus-minus with the final negative portion  $\geq 40$ ms
- 5.  $\geq$  40ms final portion of P loop upstart orthogonal X and Z leads.
- 6. Final portion of P loop delayed, notches and slurrings in the last part of the P loop
- 7. High Esophageal lead with positive P wave polarity and delayed
- 8. Low Esophageal lead with plus-minus P wave polarity and delayed
- 9. Intracavitary ECG with P wave activation craniocaudal inside the RA.
- 10. Intracavitary ECG with P wave activation caudal-cranial inside LA.

From 81,000 ECGs Bayes de Luna et al (**Bayes de Luna 1985**) collected 83 cases that fulfilled the criteria of Interatrial Conduction Disturbances with Left Atrial Retrograde Activation (IACD-LARA) (P +/- in II, III and VF with P width  $\geq$  120 msec.).

The authors present the detailed study of 35 cases with surface ECG and VCG and 29 cases with orthogonal ECG leads.

The results are then compared against two control groups: with cardiopathy (30 cases) and without cardiopathy (25 cases).

The prevalence of IACD-LARA was nearly 1% globally, and 2% among patients with valvar heart disease.

The diagnostic criteria for Interatrial Conduction Disturbances with Left Atrial Retrograde Activation (IACD-LARA) are:

- 1) ECG: P +/- in II, III and VF with P  $\ge$  120 m.
- 2) Open angle  $> 90^{\circ}$  between the first and the second part of the P.

- 3) Orthogonal ECG: P + in Y lead with a negative mode greater than 40 m.
- 4) VCG: More than 50 msec. above the X or Z axis
- 5) Duration of the P loop  $\ge 110 \text{ ms}$
- 6) Open angle between the two parts of the P loop in both frontal and right sagittal planes
- 7) Presence of notches and slurring in the last part of the P loop.

Martínez-Sellés et al studied 80 centenarians (101.4 $\pm$ 1.5 years, 21 men) with follow-ups of 6 to 34 months. Among these, 71 subjects (88.8%) underwent echocardiography. The control group comprised 269 septuagenarians. A total of 23 subjects (28.8%) had normal P wave, 16 (20%) had partial IAB, 21 (26%) had advanced IAB, and 20 (25.0%) had AF / AFI. The IAB groups exhibited premature atrial beats(PABs) more frequently than the normal P wave group also, other measurements in the IAB groups frequently fell between values observed in the normal P wave and the AF /AFI groups. These measurements included sex preponderance, mental status and dementia, perceived health status, significant mitral regurgitation, and mortality. The IAB group had a higher previous stroke rate than other groups. Compared to septuagenarians, centenarians less frequently presented a normal P wave (28.8% vs. 53.5%), and more frequently presented advanced IAB (26.3% vs. 8.2%), AF/AFI (25.0% vs. 10.0%), and PABs (28.3 vs. 7.0%) (p-values <0.01).

Relatively few centenarians (<30%) had a normal P wave, and nearly half had IAB. Particularly advanced IAB, is a pre-AF condition associated with PABs. Atrial arrhythmias and IAB occurred more frequently in centenarians than in septuagenarians. (Martínez-Sellés 2015)

## Age-related changes in P-wave morphology in healthy subjects (Havmoller 2007)

- 1) Orthogonal P-wave morphology in healthy individuals are predominantly positive in Leads X (0+ 180° in I) and Y (+90° to -90° in aVF)
- 2) In Lead Z (near the V2 line), 35% had negative morphology and 65% a biphasic one with a transition from negative to positive. The latter P-wave morphology type is significantly more common after the age of 50
- 3) P-wave increased with age being slightly longer in subjects older than 50 years old

- 4) Changes of signal averaged orthogonal P-wave morphology (biphasic signal in Lead Z), are common in healthy subjects and appear predominantly after the age of 50.
- 5) Subtle age-related prolongation of P-wave duration is unlikely to be sufficient as a sole explanation of this finding that is thought to represent interatrial conduction disturbances.

### **Atrial enlargements**

## Concept

- I. The ECG expression of atrial enlargement is due more to atrial dilation than atrial hypertrophy because the atrial wall is very thin and when submitted to an increase in pressure it usually dilates before increasing its myocardial mass (Ferroglio 1979).
- II. The standard techniques to correlate ECG changes with the presence of atrial enlargement were previously based on anatomic, radiologic, and hemodynamic standards (Reynolds 1953, Morris 1964). The necropsic studies used were feasible only in cases of very advanced heart disease, and thus their utility was limited. For more than 35 years, M-mode echocardiography and especially 2D echocardiography have been considered the "gold standard" (Reeves 1981; Kaplan 1994; Rodevan 1999) (Tables 1 and 2).

TABLE 1 Right atrial enlargement. ECG criteria with high specificity (Bayés de Luna 2013)

ECG criteria	SE %	SP %
A. QRS criteria		
1. QR or qR in V1	$\approx 15$	> 95
2. QRS V1 $\leq$ 4 mm + QRS V2/V1 $\geq$ 5	46	93
3. R/S > 1 in V1	$\approx 25$	> 95
4. $\hat{SAQRS} > 90^{\circ}$		> 95
B. P criteria		
1. P wave in inferior leads < 2.5mm	7	100
2. Positive part of P wave V1 > 1.5mm	17	100
3. Positive part of P wave $V2 > 1.5$ mm	33	100
C. Combined		
1. Positive part of P wave in V2>1.5mm+SÂQRS>+90°+R/S>1 in V1	49	100

ECG criteria		SE %	SP %
1.	Morris index (Morris 1964) (P terminal force in V1 mm/s) (0.04 mm/s)	69	93
2.	NYAC score	15	98
	P duration $\geq 120~ms$ in I or II + Morris index (>0.04 mm/s) + SÅP $\approx 0^\circ$		
3.	P duration $\geq$ 120 ms in lead II + Terminal mode negative of P in V1 > 40 ms	50	87
4.	P duration $\geq$ 120 ms in lead II + Morris index (>0.04 mm/s)	69%	49%
5.	Interpeaks of P wave >40 ms	15	100
6.	Plus-minus P-wave $\pm$ in II, III, aVF	5	100
7.	SÂP beyond $+ 30^{\circ}$	8	90
8.	P wave duration in II or III leads $\ge 0.12$ s	33%	88%

TABLE 2 Left atrial enlargement. ECG criteria based on P wave changes with high specificity

(Bayés de Luna 2013)

Cardiovascular magnetic resonance (CMR) is currently the gold standard for atrial volume assessment. It has recently been demonstrated that 2D transthoracic echocardiography consistently underestimates the left atria (LA) and right atrial (RA) volume compared with CMR imaging ( $\approx$ 15-20%) (10). However, as the volumes assessed by the two techniques have similar slopes, except for the underestimation by 2D echocardiography, the study performed using CMR to evaluate LA volume would likely result in similar results to those obtained using 2D echocardiography (**Rodevan 1999**). The accuracy of ECG criteria for left and right atrial enlargement as detected by 2D Echocardiography or CMR has recently been shown as is expressed in Tables 1 and 2. For full information consult Bayés de Luna (Bayés de Luna 2013).

The sensitivity and specificity of different criteria varies with the methodology used, and particularly with the type of population studied. The specificity is usually much higher and the sensitivity lower, but this increases if the population studied presents with a higher degree of atrial enlargement.

## Left Atrial Enlargement (LAE): Etiologies

## I) Congenital causes:

- Heart disease with pulmonary hyperflux: Ventricular Septal Defect (VSD), Persistence of Arterial Channel (PCA), Complete Atrioventricular Canal defect (CAVC defect).
- Congenital mitral stenosis

- Severe Aortic valve stenosis (AS)
- Coarctation of the Aorta
- Hypertrophic Cardiomyopathy

## II) Acquired causes:

- Mitral valvar heart disease: pure mitral valve stenosis, Mitral insufficiency/ regurgitation: "P mitrale";
- Decrease of LV compliance. E.g.: systemic hypertension, HCM, restrictive cardiomyopathies
- LVF: left ventricular failure by increase of Pd<sub>2</sub> of LV. E.g.: extensive infarctions, dilated cardiomyopathy;

## Left Atrial Enlargement (LAE) electrocardiographic criteria

## I) Direct criteria

- P wave of increased duration: ≥ 110 ms in adults, ≥ 120 ms in seniors, and 90 ms in children. Specificity: 90% and sensitivity: 40% in old age
- Notched and bifid P wave in II, with interval between the apexes ≥40 ms. Voltage of 2<sup>nd</sup> module > than the 1<sup>st</sup>. A bifid P wave is a rare finding with left atrial diameter ≥60 mm. Atrial fibrillation is present in 70% of cases.
- 3) SÂP (P axis on frontal plane) deviated to the left: between  $+40^{\circ}$  and  $-30^{\circ}$ ;
- 4) Increase in depth and duration of final negative component of the wave in V<sub>1</sub> (left atrial enlargement Morris' index (Morris 1964)); slow and deep of P in V<sub>1</sub> or V<sub>1</sub>-V<sub>2</sub>. PTFV1. P terminal force in lead V<sub>1</sub> equal or more negative than 0.04 mm/s Greater than 0.03 mm/s: product of the duration of the final negative component (duration expressed in seconds); while depth is expressed in mm. Values above 0.03 mm per second constitute a highly sensitive criterion for diagnosis of LAE.
- 5) Macruz index (Macruz 1958) > 1.7: Duration of P / Duration of PRs;
- 6) Intrinsic deflection of V1 of 30 ms (0.03 s) or greater. This deflection is measured from the apex of the initial positive component until the nadir of the final negative component of the P wave of V1.

7) P-terminal force (PTF-V1) exceeding 0.04 mm/s. This is the terminal, negative part of the P wave in lead V1 expressed as the multiplication of its depth in millimeters and width in seconds (mm/s). The normal PTF-V1 does not exceed 0.04 s wide and 1mm deep, i.e., 0.04 mm/s.

## **II) Indirect criterion**

Presence of coarse atrial fibrillation: "f" waves with amplitude equal or higher than 1 mm in  $V_1$  or  $V_1$  -  $V_2$ : in 75% of the cases of coarse AF there is LAE coexisting. There is a significant relationship between the size of the f wave and the etiology of AF. Thus, 88% of the patients with coarse AF have rheumatic valvular heart disease as underlying cause and 88% of patients with fine AF present coronary artery disease.

## False positive and false negative diagnoses of left atrial enlargement

The diagnosis of LAE may often be difficult to achieve because of the following factors:

- The presence of isolated interatrial block explains the increase in the duration of the P wave, without the presence of evident teminal negative mode of V1 (1). Thus, in contrast the combination of P wave duration in FP + increase of negative mode of P wave in V1 (Morris index) is a good criterion (Morris 1964).
- Some patients with isolated non-advanced left heart disease (e.g. mitral stenosis) without evident IAB show peaked P waves with no increase in duration (pseudo-P-pulmonale). In these cases, the presence of a P wave in V1 with a highly negative mode helps to reach a correct diagnosis (Figures below).



(A and B) P wave morphology in V1 and P loop in a case of isolated partial interatrial block (B) associated with left atrial enlargement (A)



Diagram contrasting normal and abnormal negative components of the P wave in V1. When the value calculated using the width in seconds and the height in millimetres of the negative mode exceeds 40 mm + ms, it is considered abnormal.

• If important atrial fibrosis exists, small and even unapparent P waves (concealed sinus rhythm) may be seen, even in the presence of evident left atrial or biatrial enlargement (Bayés de Luna 1978). This problem increases the number of false negatives (low SE).

Many patients with COPD or thoracic abnormalities, pectus excavatum, and those with straight-back syndrome present with short but evident negative P waves in V1, a morphology that may be confused with LAE (Bayés de Luna 2013).



#### Profile of normal P wave in RAE and LAE





Note: the findings in the Frontal Plane are not relevant for the diagnosis of LAE.

Similar to advanced interatrial block



### P loop in LAE on Horizontal plane

- The maximal vector of P is located to the left:  $\ge 0.10$  mV in adults and  $\ge 0.14$  mV in <15 years old
- Max. vector of P > 0.05 mV
- "Bow Tie" morphology



 $LA_{2/}$ : final deep and slow component:  $LAE \ge$  the area of one small square; the final minus portion indicates left atrial enlargement, abnormality or advanced IAB.

## **Right Atrial Enlargement: Etiologies**

## I) Congenital causes: "congenital P"

- I. Ebstein's anomaly "Himalayan P waves" of Taussig;
- II. Tricuspid atresia: "P tricuspidale" of Gamboa (10%);
- III. Moderate pulmonary stenosis "gothic" P (30%);
- IV. Severe pulmonary stenosis P >2.5 mm 75%;
- V. Eisenmenger Syndrome;
- VI. Atrioventricular septal defect (LV LA). RAE or BAE 60%;
- VII. Tetralogy of Fallot (T4F): only in 5% there are criteria of RAE.

## II) Acquired causes

- I. Cor pulmonale emphysema COPD: "P pulmonale"
- II. Tricuspid stenosis;
- III. Tricuspid regurgitation/insufficiency;
- IV. Double tricuspid injury;

- V. Heart failure;
- VI. Increase of RV Pd<sub>2</sub>.
- VII. Isolated Pulmonary Hypertension: the ECG is very sensitive in symptomatic patients with isolated pulmonary hypertension (Bossone 2003).

#### "Himalayan P wave"

Ebstein's anomaly is a rare, complex, fascinating, congenital anomaly with a broad pathologic-anatomical and clinical spectrum accounting for <1% of all congenital heart defects. Since its description in 1866, dramatic advances in diagnosis have been made. Very high "Himalaya mountain-like" P waves (Kaushik 2007) are observed. (The Himalayan mountain system are the planet's highest peaks around the world). P wave is >3 mm (0.3 mV) in close to 50% of cases (Armengol 1996). Tall P waves ( $\geq 2.5 \text{ mm}$ ) are attributable to right atriomegaly. A prolonged P-wave duration is occasionally registred (Jaiyesimi 1982).

In association with P wave modifications in Ebstein's anomaly, the following are frequently observed (**Blömer 1975**): Prolonged PR interval ( $\geq$ 170 ms), short PR interval if associated with WPW pattern ( $\approx$ 30% of cases), bizarre low voltage right bundle branch block pattern, initial q wave in the QRS complexes of V1 and V2 leads.

Reduced amplitude of R-wave deflections in V3R and V1, P-dextro-atriale and bizarre low RBBB without right ventricular overload almost certainly constitutes a pathognomonic finding in Ebstein's anomaly. Higher P waves and wider QRS complexes are registered in more severe cases of Ebstein's anomaly of the tricuspid valve. There is a high potential for developing arrhythmia in the vast majority of tachycardia types: atrial ectopic tachycardia, atrial flutter, atrioventricular reentry tachycardia, AV-nodal reentry tachycardia, atrial fibrillation and ventricular tachyarrhythmias.

Tendency towards multiple arrhythmogenic substrates in a single patient (Hebe 2000).

## The "Gamboa (Gamboa 1966) P wave"

The association of:

- 1. Right atrial enlargement
- 2. Diastolic, volumetric or eccentric left ventricular hypertrophy
- 3. Extreme left axis QRS deviation in the frontal plane: LAFB pattern
- 4. Counterclockwise rotation of QRS loop in the frontal plane: LAFB pattern

5. Cyanotic baby (neonate or infant). It is very suggestive of tricuspid atresia diagnosis.

From 120 children with tricuspid atresia, ECG with left axis axis deviation is observed in 94%, ERA in 58%, LVH in 96% and LAE: 47.5% (**Rosado 1987**).

The association of:

- 1. Right atrial enlargement
- 2. Diastolic, volumetric or eccentric left ventricular hypertrophy
- 3. Extreme left axis QRS deviation in the frontal plane: LAFB pattern
- 4. Counterclockwise rotation of QRS loop in the frontal plane: LAFB pattern
- 5. Cyanotic baby (neonate or infant). It is very suggestive of tricuspid atresia diagnosis.

From 120 children with tricuspid atresia, ECG with left axis axis deviation is observed in 94%, ERA in 58%, LVH in 96% and LAE: 47.5% (**Rosado 1987**).

P wave of RAE in II and V1



**Right Atrial Enlargement criteria (RAE)** 

- I) Direct ECG criteria: Direct P wave criteria are very specific but their sensitivity is very low.
- Voltage of  $P \ge 2.5$  mm in at least one of inferior leads: P pulmonale or P pulmonale parenchymal: tall and occasionally pointed P wave in inferior leads
- Aspect in apex of P wave: "Goth P"
- P wave height >1.5 mm in lead  $V_2$ . The criteria has 100% specificity preserved
- P waves with "plus-minus" pattern in right precordial leads with initial plus component ≥1.5 mm
- P wave deeply negative or positive in V<sub>1</sub>
- P wave of voltage  $\geq$  at 1.5 mm in V<sub>2</sub> in association to R/S ratio >1
- SÂP to the right of +80° (negative P wave in VL). In congenital heart diseases, SAP is not deviated to the right.
- P wave with increase in voltage and in duration in cases of extreme RAE
- Macruz index lower than  $1 = P \operatorname{duration}/_{PRs \operatorname{duration}}$  (Macruz 1958)

- The QRS criteria with a QRS amplitude in V1 <4 mm + ratio V2/V1 > 5 are highly specific (>90%) with moderate SE ( $\approx$  45 %)
- The combined P + QRS criteria with a P wave amplitude in V2 > 1.5 mm + SÂQRS
  > 90° + R/S ratio >1 in V1 in the absence of RBBB have 100% specificity and ≈ 50% sensitivity.

# II) Indirect ECG criteria

- $SAQRS_F$ : > 90°. The criteria has 100% specificity preserved
- Sodi Pallares sign<sup>1</sup>: qR, QR or qRs in  $V_1$  and  $V_2$
- Peñaloza and Tranchesi sign: QRS complexes of low voltage in V<sub>1</sub> contrasting with QRS complexes of normal voltage or increased in V<sub>2</sub>.
- R/S ratio >1 in lead V<sub>1</sub> without RBBB. The criteria has 100% specificity preserved.

## False positive and false negative diagnoses of right atrial enlargement

The ECG diagnosis of RAE may be very difficult to reach for the following reasons:

- The voltage of the P wave is strongly influenced by extra cardiac factors (Figure below), which may result in increases (hypoxia, sympathetic overdrive, etc.) (false positive) or decreases in voltage (emphysema, other barrier factors, atrial fibrosis, etc.) (false negative)
- The presence of associated fixed or intermittent atrial block may result in the transient or permanent disappearance of the ECG criteria for right atrial enlargement (false negative) (Figure in next slide)
- On the other hand, a high-voltage P wave may be seen in patients with exclusively left heart pathology and possible left atrial enlargement (false positive) (pseudo-Ppulmonale) (Bayés de Luna 2011)

These are some of the reasons why changes in the atriogram are generally not very sensitive (many false negatives) for the diagnosis of RAE. Although there are some factors that increase the incidence of false positives, they are fewer and therefore the specificity of ECG criteria for RAE is much higher.

Significant dilatation of Right Atrium: Indirect sign of RAE conditioning qR pattern in V<sub>1</sub> and V<sub>3</sub>R (Sodi-Pallares' sign) (Sodi-Pallares 1952)



Outline that explains the indirect sign of RAE: qR in V1 (sign of Sodi-Pallares). The volumetric increase of the RA, gets closer to the exploring electrode V<sub>1</sub>, recording initial QRS negativity in this lead, because this electrode records the epicardial morphology of the right atrium.

## Significance of RAE in Congestive Heart Failure (CHF)

Patients with electrocardiographic pattern of RAE, atrial flutter, AF, 1<sup>st</sup> or 2<sup>nd</sup> degree AV block Mobitz Type I, Complete LBBB and interstitial pulmonary edema with bilateral pleural effusion in chest X-rays, frequently associated to CHF( Fonseca 2004).

Atrial high resolution ECG

The duration of P wave in high resolution ECG (signal-averaged P wave (SAPWD)) significantly correlates with the size of the right atrium.(**Dixen 2004**)

## Pseudo-P-pulmonale

These cases are frequently misdiagnosed. In pure left atrial overload rarely a very high P wave ( $\geq 2.5$  mm in inferior leads) is observed with right axis deviation (P axis located to the right of +60°: negative P in VL) and major duration ( $\geq 120$  ms).

The vector of left atrial activation is directed not only posteriorly, but also more inferiorly than to the left pointing abnormally downwards.

Thus a P wave with two humps is observed: The first one, registered within the first 40 ms and responsible for a slight notch corresponds to the first part of the normal right atrial activation, while the second greater hump corresponds to the activation of the enlarged left atrium (Chow 1965, Chung 1972, Gertsch 2003).

## P-pulmonale-like patterns causes without RAE

- I. Severe hyperkalemia
- II. Asthenic habitus
- III. Enhanced sympathetic tone
- IV. Cyanosis.

## Severe hyperkalemia imitating P-pulmonale

Hyperkalemia is one of the more common acute life-threatening metabolic emergencies seen in the emergency department.

Early diagnosis and empirical treatment of hyperkalemia is dependent in many cases on the emergency physician's ability to recognize the ECG manifestations.

The ECG manifestations of hyperkalemia include (Mattu 2000):

- Diminished, flattening or absence of the P wave (sinoventricular conduction)
- Exceptionally enhanced P wave amplitude in the inferior leads imitating P pulmonale (Gertsch 2003).
- PR interval prolongation
- Widening of the QRS complexes

- A "sine-wave" appearance at severely elevated levels. The possible mechanism for the genesis of the sine wave, including loss of electrical gradient with resulting phase difference of QRS and T, associated with maintenance of His bundle activity with progressive, distal, Purkinje blockade (Sridharan 1979).
- Peaked and narrow based T waves in the precordial leads

### P loop in Right Atrial Enlargement in the Frontal Plane



P loop in Right Atrial Enlargement in the Horizontal Plane



- ➤ Maximal vector of P may exceed >0.1 mV
- $\succ$  Most of the P loop heads to the front.
- → Magnitude of anterior forces:  $\geq 0.07 \text{ mV}$
- > Max. ant. forces  $\geq 0.07$  Mv.

# P-wave components in RAE



## Biatrial Enlargement (BAE) electrocardiographic criteria

The most important diagnostic criteria are as follows

- 1. Initial component of P wave in V1 >1.5 mm and final slow and deep negative component > 1 mm in depth and 40 ms in duration
- 2. P waves of voltage >2.5 mm and duration  $\ge$ 120 ms in II
- 3. Initial part of P waves is peaked in V<sub>1</sub> and V<sub>2</sub> with voltage >1.5 mm and terminal negative mode slow (width  $\geq$ 1mm)
- 4. Signs of LAE (P waves of duration >120 ms and bimodal) with right SÂP. The opposite case is not valid because the SÂP can be on the left side in isolated RAE of patients with congenital heart diseases.
- Atrial Fibrillation associated with qR and QR type complexes in right precordial leads in the absence of myocardial infarction in V<sub>1</sub> or V<sub>1</sub>-V<sub>2</sub> (Sodi Pallares sign) (Sodi Pallares 1952)
- 6. QRS complexes of low voltage in V<sub>1</sub> contrasting with QRS complexes of normal voltage or increased in V<sub>2</sub>.

Frequently, more than one criterion is found (Figure 6) (P duration  $\ge 120$  ms in FP + P  $\pm$  in V1 with first part peaked and the second part broad and deep).

## P loop in Biatrial Enlargement in the Horizontal Plane



### Increase of anterior and posterior voltages of P loop

### Possible causes of negative P wave in lead I

- Incorrect limbs electrode cable connection during electrocardiographic recording (right to left) The frequency of ECG artifacts due to switched electrodes is 0.4% at the outpatient clinic and 4.0% at the intensive care unit (Rudiger 2007). Limb electrode misconnection: Pseudo dextrocardia by exchange of limb electrodes. All P, QRS, and T waves are negative in I, but normal progression of QRS in precordial leads rules out this hypothesis, pointing out the exchange of arm electrodes.
- Simple true dextrocardia: mirror image. Total atrio-visceral situs inversus with no heart disease. SÂP directed to the right and below, pointing at around +120 degrees (III). Negative P wave in VL and I, positive in III. Reverse progression of r wave in precordial leads V2 to V5 (decreasing).

## **Atrial T vector**

Atrial repolarization (Ta vector) is responsible for the P loop being open and not closed (0 point does not coincide with initial E point). A straight line from the onset of the P loop to the end of it, shows the magnitude and direction of the Ta vector.

#### Exteriorization of segments in vectorcardiography



In vectorcardiography, isoelectric lines corresponding to PR (PRs), ST and T-P segments are not recorded if they do not show depression or elevation. Thus, the segments manifest as stationary points. This non-manifestation is the reason why the ECG is superior to VCG in the analysis of segments and intervals.

P loop is open because atrial repolarization (TP or TA loop) is completely opposite to the P loop



In ECG, Ta or Tp wave: wave generally not visible because it is hidden by QRS. It represents atrial repolarization. Ta polarity is opposite to the P wave and its magnitude is 100 to 200 muV.

It may possibly reach the ST segment and the T wave, causing ST segment depression and resembling myocardial ischemia (Sapin 1991).

### References

1. Ariyarajah V, Spodick DH. Advanced interatrial block: a classic electrocardiogram. Cardiology. 2005;104:33-34.

- Ariyarajah V, Apiyasawat S, Puri P, Spodick DH. Specific electrocardiographic markers of P-wave morphology in interatrial block. J Electrocardiol. 2006 Oct; 39:380-384.
- 3. Ariyarajah V, Spodick DH. The Bachmann Bundle and interatrial conduction. Cardiol Rev. 2006 Jul-Aug;14:194-199.
- Armengol Rofes AJ, Serrano Durán M, Albert Brotons DC, Sánchez López C, Casaldáliga Ferrer J, Girona Comas JM. Ebstein's anomaly of the tricuspid valve. Apropos 35 cases An Esp Pediatr. 1996 Feb;44:139-144.
- 5. Asad N, Johnson VM, Spodick DH. Acute right atrial strain: regression in normal as well as abnormal P-wave amplitudes with treatment of obstructive pulmonary disease. Chest. 2003 Aug;124:560-564.
- 6. Bailin SJAtrial lead implantation in the Bachmann bundle. Heart Rhythm.2005 Jul; 2(7):784-6.
- 7. Baranchuk A, Bayés de Luna A.The P-wave morphology: what does it tell us? Herzschrittmacherther Elektrophysiol. 2015 Sep;26(3):192-199.
- 8. Herzschrittmacherther Elektrophysiol. 2015 Sep;26(3):192-199
- Bayés de Luna A, Gusí Gené C, Soler Soler J, Fort de Ribot R, Llamas Lombardia A, Roman Castillo M, TrillaSanchez E. Electrocardiologia clínica (2 volúmenes). Cinetiífico-Médica, Barcelona 1977.
- Bayes de Luna A, Fort de Ribot R, Trilla E, Julia J, Garcia J, Sadurni J, Riba J, Sagues F. Electrocardiographic and vectorcardiographic study of interatrial conduction disturbances with left atrial retrograde activation.J Electrocardiol. 1985 Jan;18:1-13.
- 11. Bayés de Luna A, Cladellas M, Oter R, Torner P, Guindo J, Martí V, Rivera I, Iturralde P.Interatrial conduction block and retrograde activation of the left atrium and paroxysmal supraventricular tachyarrhythmia. Eur Heart J. 1988 Oct; 9:1112-1118.
- 12. Bayés de Luna A. TRATADO DE ELECTROCARDIOGRAFIA CLÍNICA. Capítulo IV. Pagina 153. Editorial Cientiífico-médica. Barcelona. 1988.
- 13. Bayés de Luna A, Oter MC, Guindo J. Interatrial conduction block with retrograde activation of the left atrium and paroxysmal supraventricular tachyarrhythmias:

influence of preventive antiarrhythmic treatment. Int J Cardiol. 1989 Feb; 22:147-150.

- 14. Bayés de Luna A, Guindo J, Viñolas X, et al. Third-degree inter-atrial block and supraventricular tachyarrhythmias. Europace 1999; 1: 43
- 15. Bayés de Luna A. Clinical Arrhythmology. Wiley-Blackwell, 2011.
- Bayés de Luna A, Platonov P, Cosio FG, et al.. . Interatrial blocks. A separate entity from left atrial enlargement: a consensus report. J Electrocardiol. 2012 Sep;45(5): 445-51
- 17. Bayés de Luna A. Clinical Electrocardiography. Wiley-Blacwell 2013
- 18. Bayés de Luna A. The ECG for beginners. Wiley-Blackwell 2014.
- 19. Blömer H. Electrocardiographic and phonocardiographic findings in Ebstein's anomaly with special regard to its severity. Med Klin. 1975; 70: 1175-1178.
- Bohun CM, Potts JE, Casey BM, Sandor GG A population-based study of cardiac malformations and outcomes associated with dextrocardia. Am J Cardiol. 2007 Jul 15;100:305-309.
- 21. Bossone E, Butera G, Bodini BD, Rubenfire M. The interpretation of the electrocardiogram in patients with pulmonary hypertension: the need for clinical correlation. Ital Heart J. 2003 Dec;4:850-854.
- 22. Chavan C, Karmalkar M, Badani R, et al. Evaluation of bachmann bundle pacing versus right atrial pacing in prevention of atrial fibrillation after coronary artery bypass surgery. Indian Pacing Electrophysiol J. 2011 Feb 7;10(12):529-35.
- 23. Chou TC, Helm RA. The Pseudo P Pulmonale. Circulation. 1965; 32: 96-105.
- 24. Chung DK, Chung EK. Pseudo-P-pulmonale. W V Med J. 1972; 68:10-11.
- 25. Chung EK. Aberrant atrial conduction. Unrecognized electrocardiographic entity. Br Heart J. 1972 Apr;34(4):341-6.
- 26. Conde D, Baranchuk A. Interatrial block as anatomical-electrical substrate for supraventricular arrhythmias: Bayeś syndrome. Arch Cardiol Mex.2014 Jan-Mar; 84(1):32-40.

- Conde D, Baranchuk A, Bayés de Luna A.Advanced interatrial block as a substrate of supraventricular tachyarrhythmias: a well recognized syndrome.J Electrocardiol. 2015 Mar-Apr;48(2):135-40.
- 28. Daubert C, Gras D, Berder V et al [Permanent atrial resynchronization by synchronous bi-atrial pacing in the preventive treatment of atrial flutter associated with high degree interatrial block]. Arch Mal Coeur Vaiss. 1994 Nov;87(11 Suppl): 1535-46.
- 29. Dixen U, Joens C, Rasmussen BV, Parner J, Jensen GB. Signal-averaged P wave duration and the dimensions of the atria. Ann Noninvasive Electrocardiol. 2004 Oct; 9:309-315.
- 30. Ferroglio S, Pham TD, Hordof A, et al. Right atrial ultrastructure in congenital heart disease. Am J Cardiol 1979; 43: 820.
- 31. Fonseca C, Oliveira AG, Mota T, Matias F, Morais H, Costa C, et al; Evaluation of the performance and concordance of clinical questionnaires for the diagnosis of heart failure in primary care. EPICA Investigators.Eur J Heart Fail. 2004 Oct;6(6): 813-20, 821-2.
- 32. Gamboa R, Gersony WM, Nadas AS. The electrocardiogram in tricuspid atresia and pulmonary atresia with intact ventricular septum. Circulation. 1966 Jul;34(1):24-37.
- Gertsch Marc The ECG A two-Step Approach to Diagnosis 2003; Chapter 4 pp: 48 and 52.
- Havmoller R, Carlson J, Holmqvist F, Herreros A, Meurling CJ, Olsson B, Platonov P. Age-related changes in P wave morphology in healthy subjects. BMC Cardiovasc Disord. 2007 Jul 27;7:22.
- 35. Hebe J. Ebstein's anomaly in adults. Arrhythmias: diagnosis and therapeutic approach. Thorac Cardiovasc Surg. 2000; 48: 214-219.
- 36. Holmqvist F, Platonov PG, Carlson J, et al. Variable interatrial conduction illustrated in a hypertrophic cardiomyopathy population. Ann Noninvasive Electrocardiol. 2007 Jul;12(3):227-36.
- 37. Jairath UC, Spodick DH. Exceptional prevalence of interatrial block in a general hospital population. Clin Cardiol. 2001 Aug;24(8):548-50.
- 38. Jaiyesimi F.Observations on the so-called non-specific electrocardiographic changes in endomyocardial fibrosis. East Afr Med J. 1982 Jan;59:56-69.

- 39. Julia J, Bayes De Luna A, Candell J, et al. Auricular aberration: apropos of 21 cases. Rev Esp Cardiol. 1978;31(2):207-14.
- 40. Kaplan J, Evans G, Foster E, et al. Evaluation of ECG criteria for right atrial enlargement by quantitative two-dimensional echocardiography. J Am Coll Cardiol 1994; 23: 747.
- 41. Kaushik ML, Sharma M, Kashyap R. 'Himalayan' p wave.J Assoc Physicians India. 2007;55: 856. Kitkungvan D, Spodick DH. Interatrial block: is it time for more attention? J Electrocardiol. 2009 Nov-Dec;42:687-692Kitkungvan D, Spodick DH. Interatrial block: is it time for more attention? J Electrocardiol. 2009 Nov-Dec; 42:687-692.
- 42. Lee KS, Appleton CP, Lester SJ,et al. Relation of electrocardiographic criteria for left atrial enlargement to two-dimensional echocardiographic left atrial volume measurements. Am J Cardiol. 2007 Jan 1;99(1):113-8.
- 43. Macruz R, Perloff JK, Case RB. A method for the electrocardiographic recognition of atrial enlargement. Circulation. 1958 May;17:882-889.
- 44. Martínez-Sellés M, Roessel AM, Álvarez-García J, de la Villa BG, Cruz-Jentoft AJ, Vidán MT, Díaz JL, Felix Redondo FJ, Durán Guerrero JM, Bayes-Genis A, de Luna AB; investigators of the Cardiac and Clinical Characterization of Centenarians (4C) registry.Interatrial Block and Atrial Arrhythmias in Centenarians: Prevalence, Associations, and Clinical Implications Heart Rhythm. 2015 Oct 28. pii: S1547-5271(15)01364-8.
- 45. Mattu A, Brady WJ, Robinson DA. Electrocardiographic manifestations of hyperkalemia. Am J Emerg Med. 2000;18:721-729.
- 46. Mehrzad R, Spodick DH. Interatrial block: a virtual pandemic requiring attention. Iran J Med Sci. 2014 Mar;39(2):84-93.
- 47. Morris JJ Jr, Estes EH Jr, Whalen RE, Thompson HK Jr, Mcintosh HD. P-WAVE ANALYSIS IN VALVULAR HEART DISEASE. Circulation. 1964 Feb;29:242-252
- 48. Platonov PG. Atrial conduction and atrial fibrillation: what can we learn from surface ECG? Cardiol J. 2008;15:402-407
- 49. Reeves WC, Hallahn W, Schwitter EEJ, et al. Two-dimensional echocardiographic assessment of ECG criteria for right atrial enlargement. Circulation 1981; 64: 387.
- 50. Reynolds G. The atrial electrogram in mitral stenosis. Br Heart J 1953; 15: 250.

- 51. Rodevan D, Bjornerheim R, Ljosland M, et al. Left atrial volumes assessed by three-and two- dimensional echocardiography compared to MRI estimates. Int J Cardiol Imaging 1999; 15: 397.
- 52. Rosado-Buzzo AA, Santamaría-Díaz H, Gómez-Gómez M, Alva-Espinosa C, Maulen-Radovan X, Palacios-Macedo X. Tricuspid atresia. Clinical course in 120 childrenArch Inst Cardiol Mex. 1987 Sep-Oct;57:375-381.
- 53. Rudiger A, Hellermann JP, Mukherjee R, Follath F, Turina J. Electrocardiographic artifacts due to electrode misplacement and their frequency in different clinical settings. Am J Emerg Med. 2007 Feb;25(2):174-8.
- 54. Sapin PM, Koch G, Blauwet MB, McCarthy JJ, Hinds SW, GettesLS. Identification of false positive exercise tests with use of electrocardiographic criteria: a possible role for atrial repolarization waves. J Am Coll Cardiol. 1991; 18: 127-135.
- 55. Sodi Pallares D, Bisteni A, Hermann GR. Some views on the significance of qR and QR type complexes in right precordial leads in the absence of myocardial infarction. Am Heart J 1952;43:716-734.
- 56. Spodick DH. Unappreciated prevalence of interatrial block and associated consequences: a poorly perceived pandemic.Mayo Clin Proc. 2004 May;79(5): 668-70.
- 57. Spodick DH, Ariyarajah V, Apiyasawat S. Higher prevalence of cardiovascular events among patients with abnormal atrial depolarization and coronary artery disease at 18 months' post-exercise tolerance testing. Am Heart Hosp J. 2007 Fall; 5(4):236-40
- 58. Spodick DH, Ariyarajah V. Interatrial block: a prevalent, widely heglected and portentous abnormality. J Electrocardiol 2008; 41: 61.
- 59. Spodick DH, Ariyarajah V.Interatrial block: the pandemic remains poorly perceived. Pacing Clin Electrophysiol. 2009 May;32:667-672.
- 60. Sridharan MR, Horan LG, Flowers NC Combined effects of graded hyperkalemia on activation and recovery. Am Heart J. 1979;97:622-630.
- 61. Tsao CW, Josephson ME, Hauser TH, et al. Accuracy of electrocardiographic criteria for atrial enlargement: validation with cardiovascular magnetic resonance. J Cardiovasc Magn Reson. 2008 Jan 25;10:7.

62. Waldo A, Harry L, Bush Jr, et al. Effects on the canine P wave of discrete lesions in the specialized atrial tracts. Circulation Res 1971; 29: 452.