

# **ECG restitution of the beat-to-beat QT-TQ intervals in humans. Can we begin assess arrhythmia vulnerability?**


Anthony A. Fossa, PhD.

Research Fellow

Dept of Translational and Molecular Medicine,  
Pfizer Inc.



# Conceptualized view of QT-RR relationship




# ECG Restitution

- Ability of the heart to recover from one beat to the next.
- Examines the relationship between action potential duration (QT interval) and diastolic interval (TQ interval) through the ECG.
- Replaces the traditional invasive electrophysiology procedure
- As QT/TQ ratio increases between beats this may be associated with increase arrhythmia vulnerability due to re-entry




# Effects of heart rate on restitution and alternans



Am J Physiol 275 (Heart Circ Physiol. 44): H1635-1642, 1998



# QT prolongation during tachycardia dramatically impairs restitution (increase QT/TQ ratio >1)



## Methods in normals and TdP

- Sequential beat-to-beat analyses of QT, TQ, and RR
- 22.5-hour Holter on 3 successive days
- Baseline, 160 (n= 38) or 320 (n=19) mg oral Sotalol

### TdP Case Study



- 66 yo female CAD: Holter after sotalol (2 mg/kg, iv) prior to TdP

## Restitution Parameters

- **Lower TQ 5<sup>th</sup> quantile:** boundary of lower 5% of beats
- **% Beats with QT/TQ > 1:** Reflects relative time spent in on steep portion of restitution relationship
- **Upper 98% quantile of QT/TQ ratio:** reflects degree and magnitude of steepness in restitution relationship for beats that may pose the greatest risk





# Baseline in normals volunteers



Parameters	Baseline Assessment Periods (98% Confidence Bounds)		
	22.5 Hours	Cmax (Hours 2-4 )	Nocturnal (Hours 19-21)
RR interval (ms)	828 (794-858)	827 (794-866)	994 (947-1050) #
QT interval (ms)	375 (366-383)	373 (364-382)	411 (401-420) #
QTc interval (Bazett)	412 (405-420)	409 (401-418)	413 (405-422)
TQ interval (ms)	456 (422-481)	456 (425-488)	586 (538-629) #
TQmin <sub>5th quantile</sub> (ms)	270 (252-287)	298 (279-321)	404 (369-442) #
% (QT/TQ ratio) > 1	25 (18-34)	20 (13-32)	6 (4-9) #
(QT/TQ ratio) <sub>max 98% quantile</sub>	1.52 (1.43-1.62)	1.41 (1.31-1.50)	1.23 (1.13-1.34) #



# Sotalol improves restitution despite QT prolongation





## Comparison of Cmax period to time-matched baseline

**RR:** ↑ 235 ms  
**QT:** ↑ 101 ms  
**TQ:** ↑ 134 ms  
**QTc:** ↑ 52 ms

**TQ 5th Quantile:** ↑ 115 ms  
**QT/TQ 98th Quant:** ↓ 15%  
**%QT/TQ beats >1:** ↓ 30%



# Restitution is impaired prior to TdP



## Comparison to Cmax period from normals

RR: ↓ 172 ms  
QT: ↑ 53 ms  
TQ: ↓ 225 ms  
QTc: ↑ 95 ms

TQ 5th Quantile: ↓ 126 ms  
QT/TQ 98th Quant: ↑ 58%  
%QT/TQ beats >1: ↑ 722%



# ECG Restitution Summary

- Can be obtained through digitized Holter recordings
- May be used to assess changes in autonomic state in conjunction with QT prolongation
- If boundaries can be defined, could delineate normal from abnormal repolarization status.



# Other Contributors

- Pfizer
  - Todd Wisialowski
  - Eric Wolfgang
  - Kim Crimin
- Daiichi-Sankyo
  - Nenad Sarapa
- AMPS
  - Fabio Badilini
- U. Rochester Med. Ctr.
  - Jean-Philippe Couderc
  - Wojciech Zareba
- LMU Munchen,  
Germany
  - Stefan Kaab
  - Martin Hinterseer

