


Brugada Phenocopies

Andrés Ricardo **Pérez-Riera**, MD PhD
Raimundo **Barbosa-Barros**, MD
Rodrigo **Daminello-Raimundo**, PhD
Luiz Carlos **de Abreu**, PhD
Kjell **Nikus**, MD PhD



First Edition

International Registry on Brugada Phenocopy

www.brugadaphenocopy.com

Brugada phenocopy

“ A phenotypic trait or disease that resembles the trait expressed by a particular genotype, but in an individual who is not a carrier of that genotype”.

Phenocopy

A variation in an organism that resembles a genetic one, but has an environmental rather than a genetic cause, and is not inherited

The definition of the Brugada syndrome: Point of view

Brugada syndrome (BrS) is a hereditary arrhythmic entity with a low and incomplete worldwide prevalence (0.5 per 1,000 or 5 to 20 per 10,000 individuals), predominant in Southeast Asia (prevalence of 3.7 per 1,000) and in general much more frequent in Asians in relation to other ethnic groups: nine times more common in relation to Caucasians, 36 times more common than in Hispanics and almost non-existent in the north of the African continent Denmark. The entity is also highly prevalent in males (male / female ratio 9: 1 in Southeast Asia and 3: 1 male/female ratio in Caucasians), caused by alterations in the structure and function of certain cardiac ion channels and less expression of connexin 43 (Cx43) in the right ventricle predominantly in ventricular outflow tract (RVOT) causing electromechanical abnormalities. The lower and heterogeneous expression of Cx43 produces functionally significant electrophysiological heterogeneity in the thickness of the ventricular wall and may be a mechanism to promote transmural dispersion of repolarization. Until recently, it was considered an autosomal dominant Mendelian entity in $\approx 25\%$ of cases or sporadic, although it is currently thought to be most likely an oligogenic disorder, rather than a Mendelian condition *, affecting several loci, influenced by environmental factors and whose Diagnosis is based on the presence of a spontaneous or drug-induced coved-type ST segment elevation characterized by elevation of the J point and the ST segment of ≥ 2 mm, of superior convexity “coved type” (Subtype 1A) or descending rectilinear (Subtype 1A) followed by a symmetric negative T wave in ≥ 1 high right and / or right precordial lead; with (Subtype 1C) or without associated early repolarization pattern in the inferior or inferolateral wall with an increased risk of syncope, (fainting), palpitations, precordial pain, seizures, difficulty breathing (nocturnal agonal respiration), and / or sudden cardiac death (SCD) secondary to polymorphic ventricular tachycardia / ventricular fibrillation (PVT/VF) unexplained cardiac arrest or documented VF/polymorphic VT or paroxysmal atrial fibrillation in the absence of **macroscopic** or **apparent** structural heart disease, electrolyte disturbance, use of certain drugs or coronary heart and fever. and that typically occurs midnight-to-early-morning period at rest ($\approx 80\%$ of cases) or or at a low level of physical activity especially during sleep, which suggests

that the parasympathetic is a determining factor in, arrhythmogenesis: Higher level of vagal tone and higher levels of Ito at the slower heart rates. Although it is considered a genetic disease, its mechanism remains unknown in $\approx 70-75\%$ of cases and a single mutation is not sufficient to cause the BrS phenotype cases. Although $\approx 20\%$ of patients with BrS carry mutations in *SCN5A*, the molecular mechanisms underlying this condition are still largely unknown.. BrS is associated in with mutations in the *SCN5A* gene, which encodes for the pore-forming α subunit of the cardiac Na^+ channels. *SCN5A*, which was identified as the first BrS-associated gene in 1998, has emerged as the most common gene associated with BrS, Only the *SCN5A* gene is classified as having definitive evidence as a cause for BrS. The role of genetics in the approach to the arrhythmic patient, progressing beyond the concept of “one mutation, one disease”, and raising concerns about the most appropriate approach to patients affected by structural/electrical cardiomyopathy. Currently, the best model is the human patient population and probably BrS is an oligogenic disease (1) There have been more than 400 mutations in *SCN5A* gene that have been associated with BrS. This evidence-based review of genes reported to cause (BrS) and routinely clinically tested in patients indicates that 20 of 21 genes lack sufficient genetic evidence to support their causality for BrS. Type 2 pattern has also been associated with mutations in *SCN5A*, glycerol-3-phosphate dehydrogenase 1-like (*GPD1L*), which is the domain responsible for a site homologous to *SCN5A*, and *CACNA1C*, the gene responsible for the α -subunit of cardiac L-type calcium channels (LTCC) Mutations of 21 genes other than *SCN5A* have been implicated in the pathogenesis of BrS to date. Multiple pathogenic variants of genes have been shown to alter the normal function of Na^+ , K^+ Gain-of-function mutations in genes encoding for potassium channels have also been implicated in BrS. Genes influencing Ito include *KCNE3*, *KCND3* and *SEMA3A* (semaphoring, an endogenous K^+ channel inhibitor) while *KCNJ8* and *ABCC9* (encoding for *SUR2A*, the ATP-binding cassette transporter for the KATP channel) mutations affected the IK_{ATP} .

1. Michelle M. et al. Brugada Syndrome: Oligogenic or Mendelian Disease? Int J Mol Sci. 2020 Mar; 21(5): 1687. Published online 2020 Mar 1. doi: 10.3390/ijms21051687

KCNH2, which encodes for IKr was proposed by Wang et al.(1) to be involved in BrS development.

Dysfunction in the KCNAB2, which encodes the voltage-gated K⁺ channel β2-subunit, was associated with increased Ito activity and identified as a putative gene involved in BrS. Kvβ2 dysfunction can contribute to the Brugada electrocardiographic pattern.(2)

Classification of hereditary diseases

○ **Monogenic or Mendelian**, to be transmitted to the offspring according to Mendel's laws. They can be

1. Autosomal Dominant,
2. Autosomal Recessive, or
3. X-linked.

Mendelian inheritance refers to the patterns of inheritance that are characteristic of organisms that reproduce sexually. It refers to the type of inheritance that can be easily understood as a consequence of a single gene.

○ **Multifactorial or polygenic**: produced by mutations in several genes, generally of different chromosomes and the combination of multiple environmental factors (age, sex, bad habits (obesity, tobacco, alcohol), toxic environments or a limited childhood.) And

○ **Polygenic Oligogenic***, where there are a few genes that have more influence than the rest. In the case of BrS, the SCN5A gene and that depending on its presence other mutations are expressed (epistasis: <https://academic.oup.com/hmg/article/11/20/2463/616080>). Despite their importance, mutations in the SCN5A gene are present in ≈20 to 30% of cases.

1. **Q Wang 1, Seiko Ohno , Wei-Guang Ding , Megumi Fukuyama , Akashi Miyamoto, Hideki Itoh , Takeru Makiyama ,et al Gain-of-Function KCNH2 Mutations in Patients with Brugada Syndrome. J Cardiovasc Electrophysiol . 2014 May;25(5):522-530. doi: 10.1111/jce.12361.**

2. **Vincent Portero, Solena Le Scouarnec, Zeineb Es-Salah-Lamoureux, Sophie Burel, Jean-Baptiste Gourraud, Stéphanie Bonnaud, Pierre Lindenbaum, et al. Dysfunction of the Voltage-Gated K⁺ Channel β2 Subunit in a Familial Case of Brugada Syndrome,J Am Heart Assoc. 2016 Jun; 5(6): e003122. doi: 10.1161/JAHA.115.003122**

Variations to Mendel's Laws Extensions and Exceptions

In these cases, genotypic ratio is a Mendel predicted but phenotypic ratio is altered

- 1) Lethal allele combinations** A pair of identical alleles that are both present in an organism that ultimately results in death of that organism are referred to as recessive lethal alleles. Though recessive lethals may code for dominant or recessive traits, they are only fatal in the homozygous condition. Examples of diseases caused by recessive lethal alleles are cystic fibrosis, Tay-Sachs disease, sickle-cell anemia, and **brachydactyly**.
- 2) Multiple alleles:** Today, we know that not all alleles behave quite as straightforwardly as in Mendel's experiments. For example, in real life: Allele pairs may have a variety of dominance relationships (that is, one allele of the pair may not completely "hide" the other in the heterozygote). There are often many different alleles of a gene in a population.
- 3) Different dominance relationships** Dominance is a relationship between two alleles of a gene and their associated phenotypes. A "dominant" allele is dominant to a particular allele of the same gene that can be inferred from the context, but it may be recessive to a third allele, and codominant to a fourth.
- 4) Epistasis** is a phenomenon in genetics in which the effect of a gene mutation is dependent on the presence or absence of mutations in one or more other genes, respectively termed modifier genes. In other words, the effect of the mutation is dependent on the genetic background in which it appears. Epistatic mutations therefore have different effects on their own than when they occur together. Originally, the term *epistasis* specifically meant that the effect of a gene variant is masked by that of a different gene.

4. **Penetrance and expressivity:** is used to describe whether or not there is a clinical expression of the genotype in the individual. Expressivity is the term that describes the differences observed in the clinical phenotype between two individuals with the same genotype.
5. **Pleiotropy** occurs when one gene influences two or more seemingly unrelated phenotypic traits. Such a gene that exhibits multiple phenotypic expression is called a pleiotropic gene. Mutation in a pleiotropic gene may have an effect on several traits simultaneously, due to the gene coding for a product used by a myriad of cells or different targets that have the same signaling function.
6. **Genetic heterogeneity:** Genetic heterogeneity occurs through the production of single or similar phenotypes through different genetic mechanisms. There are two types of genetic heterogeneity: allelic heterogeneity, which occurs when a similar phenotype is produced by different alleles within the same gene; and locus heterogeneity, which occurs when a similar phenotype is produced by mutations at different loci
7. **Phenocopies** The term was coined by Richard Goldschmidt in 1935. He used it to refer to forms, produced by some experimental procedure, whose appearance duplicates or copies the phenotype of some mutant or combination of mutants. A phenotypic trait or disease that resembles the trait expressed by a particular genotype, but in an individual who is not a carrier of that genotype

Definition and Diagnostic Criteria for Brugada Phenocopy(1)

Brugada Phenocopies(BrPs) are clinical entities characterized by Brugada ECG pattern type I and type II in the absence of BrS. A phenotypic trait or disease that resembles the trait expressed by a particular genotype, but in an individual who is not a carrier of that genotype. an environmentally induced, nongenetic alteration of a phenotype to a form that resembles the expression of a known genetic mutation.

Diagnostic criteria for BrP are:

- Characteristic type I or type II Brugada ECG pattern
- The presence of an underlying condition inducing Brugada ECG patterns
- Resolution of the ECG patterns after eliminating the underlying condition
- Low pretest probability for BrS as defined by lack of clinical symptoms, medical history and family history suggestive of BrS
- Negative provocative testing with sodium channel blocker drugs (not mandatory if surgical RVOT manipulation has occurred within the last 96 hours)
- A negative genetic testing for SCN5A (recommended but not mandatory as it is possible to identify mutation only in 20%-30% of probands that are known to have BrS.

1. Daniel D Anselm, Jennifer M Evans, and Adrian Baranchuk, Brugada phenocopy: A new electrocardiogram phenomenon World J Cardiol. 2014 Mar 26; 6(3): 81–86. doi: 10.4330/wjc.v6.i3.81

Diagnostic criteria for Brugada phenocopy (First 4 criteria are mandatory)

1. ECG pattern has type 1 or type 2 Brugada morphologic criteria
2. Presence of an underlying condition that is identifiable and reversible
3. Resolution of the ECG pattern upon elimination of the underlying condition
4. Low pretest probability for Brugada syndrome determined by the lack of symptoms, clinical history, and family history
5. A negative provocative test with a sodium channel blocker drug (e.g., ajmaline, flecainide, or procainamide)*
6. A negative genetic test**

*Provocative testing is not mandatory if surgical right ventricular outflow tract manipulation has occurred within the last 96 hours.

**Desirable, but not mandatory, since the SCN5A mutation has been identified in only 20-30% of probands affected by Brugada syndrome.

ECG: Electrocardiogram.

The characteristic ST-segment elevation seen in BrPps can be explained by a transmural gradient that arises from an accentuated I_{to} -mediated action potential notch and a loss of the AP dome in the epicardium but not the endocardium. The loss of the AP dome can result from a disruption in the homeostasis of active inward and outward currents at the end of phase 1 of the AP. Specifically, any mechanism that increases (\uparrow) outward, I_{to} , adenosine triphosphate-sensitive potassium current (I_{K-ATP}), delayed rectifier potassium current [I_{Ks}, I_{Kr}] or decreases outwards currents (i.e., I_{Ca-L} , fast I_{Na}) will result in the characteristic ST-segment elevation seen in the Brugada Type-1 ECG. This provides a possible explanation of the general, underlying pathogenesis of the diverse forms of BrPps. (**Shimizu W. Acquired forms of the Brugada syndrome. *J Electrocardiol* 2005;38:22–25.**)

Morphologic classification system from the International Registry of Brugada Phenocopies

Type	
Type 1	BrP with a coved Brugada ECG pattern
Type 2	BrP with a saddleback Brugada ECG pattern
Class	
Class A sodium	All mandatory BrP diagnostic criteria are satisfied, including positive results from a provocative drug challenge with channel blockers
Class B	Highly suspected BrP: Not all mandatory criteria are complete*
Class C	Highly suspected BrP: Provocative drug challenge is not indicated and/or justified**

*This group includes cases in which a provocative drug challenge is pending or it is not possible to perform due to various factors, such as the patient is lost to follow-up or deceased.

**This group includes cases in which a provocative drug challenge is not clinically indicated, such as patients with recent surgical right ventricular outflow tract manipulation or BrP secondary to the use of inappropriate high-pass ECG filtering. BrP: Brugada phenocopy; ECG: Electrocardiogram.

Etiologic classification for Brugada phenocopy

- 1. Metabolic conditions: Hyperkalemia and severe hypokalemia due to gastrointestinal loss.(1;2;3)**, hyponatremia, hypercalcemia, hypothermia
- 2. Mechanical compression** with involvement of the RVOT by either mediastinal mass, intracardiac tumors or metastatic carcinomas. .(4)
- 3. Ischemia and pulmonary embolism** ischemia-induced transient ion channel dysfunction generating \uparrow Ito current and decreased \downarrow Na+.(5)
- 4. Myocardial and pericardial disease:** Chagas myopathy,(6;7) acute and chronic myocarditis,(8) acute pericarditis(9)
- 5. Electrocardiogram modulation**
- 6. Miscellaneous:** consumption of yellow oleander seeds (T. Peruviana). (10) Methanol Intoxication (11)

- 1. Xu G, Gottschalk BH, Anselm DD, Benditt DG, Maheshwari A, Sreenivasan S, et al. Relation of the Brugada phenocopy to hyperkalemia (from the International Registry on Brugada Phenocopy). Am J Cardiol 2018;121:715–7.**
- 2. Genaro NR, Anselm DD, Cervino N, Estevez AO, Perona C, Villamil AM, et al. Brugada phenocopy clinical reproducibility demonstrated by recurrent hypokalemia. Ann Noninvasive Electrocardiol 2014;19:387–90.**
- 3. Neto NO, Pilla L, Oliveira WS, Velloso RN, Maia FG, do Nascimento AL, et al. Brugada Phenocopy induced by recurrent hyperkalemia: More evidence for the reproducibility of a new phenomenon. J Electrocardiol 2018;51:402–4.**
- 4. Pérez-Riera AR, Barbosa Barros R, Daminello-Raimundo R, Resende Barbosa MPC, de Abreu LC. Brugada phenocopy caused by a compressive mediastinal tumor. Ann Noninvasive Electrocardiol 2018;23:e12509.**
- 5. Peters S. Brugada phenocopy in percutaneous coronary intervention of the right coronary artery. Int J Cardiol 2016;203:675.**
- 6. Arce M, Riera AR, Femenía F, Baranchuk A. Brugada electrocardiographic phenocopy in a patient with chronic Chagasic cardiomyopathy. Cardiol J 2010;17:525–7. 49.**
- 7. Brito MR, Miranda CE, Rabelo W, Marino RL. Type 1 electrocardiographic Brugada pattern in a woman with Chagas disease: a case report. Europace 2010;12:1345–6.**
- 8. Li A, Tung R, Shivkumar K, Bradfield JS. Brugada syndrome— Malignant phenotype associated with acute cardiac inflammation? HeartRhythm Case Rep 2017;3:384–8.**
- 9. Yu M, Zhang Q, Huang X, Zhao X. Type 1 Brugada phenocopy in a patient with acute pericarditis. J Electrocardiol 2018;51:1121–3.**
- 10. R Gunaseelan 1, M Sasikumar 2, K Aswin 2, Shirshendu Dhar 2, N Balamurugan 2, Vivekanandan Pillai 2Brugada phenocopy induced by consumption of yellow oleander seeds - A case report. J Electrocardiol. 2020 Jul 24;62:107-109. doi: 10.1016/j.jelectrocard.2020.07.014. Online ahead of print. J Electrocardiol. 2020 Jul 24;62:107-109. doi: 10.1016/j.jelectrocard.2020.07.014**
- 11. Jesús Monterrubio-Villar 1, David Llinares-Moya 1Brugada Phenocopy Induced by a Lethal Methanol Intoxication Eur J Case Rep Intern Med. 2020 Jan 1;7(2):001374. doi: 10.12890/2020_001374**

ECG modulation

Another underlying condition for BrP is the improper application of high-pass filtering. Normally a high-pass filter is used for eliminating low frequency noises during ECG recording which are most likely resulted from movement of ECG electrodes on the skin, physicochemical skin-electrode changes and body movement related to normal breathing. Low frequency components of normal ECG such as ST segment can be distorted in case applying non-recommended cutoff frequency values. Importantly, although the change in the ST segment by applying improper high-pass filter depends on the baseline ECG characteristics, it is more common in the right precordial leads. When application of a nonstandard high-pass filter of 0.5 Hz produced Brugada ECG pattern. (1)

Miscellaneous

BrP in patients with Ebstein anomaly, (2) intracranial hemorrhage (3) and following biphasic synchronized electrical cardioversion for atrial fibrillation. (4)

References

- García-Niebla J, Serra-Autonell G, Bayés de Luna A. Brugada syndrome electrocardiographic pattern as a result of improper application of a high pass filter. Am J Cardiol 2012;110:318– 20.**
- 1. Kaiser E, Pastore CA, Moffa PJ. Ebstein's anomaly with Brugada-like electrocardiogram pattern: A critical view. Cardiol J 2010;17:415. 55.**
 - 2. Labadet C, Gottschalk BH, Rivero M, Hadid C, Fuselli J, Anselm DD, et al. Brugada phenocopy in the context of intracranial hemorrhage. Int J Cardiol 2014;177:e156–7.**
 - 3. Abdalla KA, Sorgente A, Brugada P. Transient appearance of Brugada-like Type 1 electrocardiogram pattern immediately after biphasic synchronized electrical cardioversion for atrial fibrillation: a case series. European Heart Journal-Case Reports 2019;1–8.**

Recommendations for future Brugada phenocopy case report publication (1)

1. It is mandatory to include a 12-lead ECG tracing with emphasis on the right precordial leads V_1-V_3 .
2. The inclusion of additional leads is desirable and highly recommendable.(2)
3. It is paramount to comment on the presence of a past medical history of syncope and a family history of sudden death or syncope, which can be of help in differentiating between BrS from BrPs. Including whether or not provocative testing, genetic testing
4. To perform an ECG tracings of immediate family members were recorded is also important in helping to rule out true BrS.
5. To report whether the Brugada ECG pattern resolved after the treatment of the underlying cause; otherwise it is difficult to classify them as a BrP, because we cannot know whether the underlying condition was truly behind the Brugada ECG manifestation.
6. It is important to document the resolution of the Brugada ECG to infer a direct association between the “environmental factor” and the Brugada ECG pattern.
7. The transient nature of the ECG alteration is a hallmark in differentiating Brugada syndrome from Brugada phenocopy(3)


1. **Adrian Baranchuk 1, Timothy Nguyen, Min Hyung Ryu, Francisco Femenía, Wojciech Zareba, Arthur A M Wilde, Wataru Shimizu, Pedro Brugada, Andrés R Pérez-Riera. Brugada phenocopy: new terminology and proposed classification, *Ann Noninvasive Electrocardiol.* 2012 Oct;17(4):299-314. doi: 10.1111/j.1542-474X.2012.00525.x.**
2. **Teijeiro R, Garro HA, Acunzo RS, *et al.* Recording of high V1-V3 precordial leads may be essential to the diagnosis of Brugada syndrome during the ajmaline test. *J Cardiovasc Pharmacol Ther* 2006;11:153–155.**
3. **Lukyanetz EA, Shkryl VM, Kostyuk PG. Selective blockade of N-type calcium channels by levetiracetam. *Epilepsia.* 2002;43(1):9–18**

Using simulations of the anterior ventricular wall Rivera –Juarez et al. showed that potassium concentration, fibrosis, and I_{to} are all involved the development of the Brugada pattern and act synergistically to shape the characteristic pattern (**Rivera-Juárez A, Hernández-Romero I, Puertas C, et al. Clinical characteristics and electrophysiological mechanisms underlying Brugada ECG in patients with severe hyperkalemia. J Am Heart Assoc. 2019;8(3):e010115.**)

Brugada phenocopy examples



Brugada phenocopy caused by a compressive mediastinal tumor

We presented a case of mediastinal non-Hodgkin's lymphoma that compressed the RVOT originating a type 1 Brugada ECG pattern. After the treatment and lymphoma reversal, this pattern disappeared. The negative provocative ajmaline test indicates low probability of Brugada syndrome.




Initial ECG. STE in aVR and from V_1 – V_3 decreasing progressively (this characteristic differentiates RV infarction from anteroseptal LV infarction). Concomitantly, ST depression is observed in inferior and lateral walls. Conclusion: type 1 Brugada ECG pattern


Posteroanterior chest x-ray showed a large pleural effusion on the right side that disappeared after treatment.




Posteroanterior chest x-ray. Large pleural effusion before (a) and after treatment (b)



The computed tomography indicates the tumor extension (arrow)




ECG normalization after oncologic therapy. Low QRS voltage in the limb leads (<5 mm) and precordial leads (<10 mm), ST depression in inferior and lateral precordial leads



Clinical diagnosis: terminal renal insufficiency. Severe hyperkalemia: K^+ 8.7 mEq/L. This sign is known as dialyzable injury current. ECG diagnosis: very likely, junctional with P waves near J point, HR: 54 bpm, QRSd: 160 ms, ST segment elevation from V_1 to V_3 and I, aVL and aVR. V_1 to V_3 display ST segment upwardly convex pattern, similar to Brugada syndrome or Brugada phenocopy”, typical T waves in “tent”, pointed, and with a narrow base.

Next figure shows a Brugada phenocopy secondary to accidental plasma concentrations of propafenone in the toxic range.



ECG diagnosis: Left atrial enlargement, PR interval prolongation or first-degree AV block secondary to augmentation of effective refractory periods of atrioventricular node ($>$ AH interval), His-Purkinje system ($>$ HV interval), nonspecific intraventricular conduction disturbance, (marked abnormal ventricular activation pattern (bizarre): does not satisfy the criteria of either LBBB or RBBB), long QT interval with normal JT interval and Brugada type 1 ECG phenocopy: ST segment elevation convex to the top followed by negative T waves from V_1 to V_3 Induced Brugada-type 1 ECG pattern, is a sign for imminent malignant arrhythmias.

Brugada Phenocopy References

- Abraham C, Maharaj S. DKA-induced Brugada phenocopy mimicking STEMI. *Heart Asia*. 2018 May 26;10(2):e011027.
- Abusin S. Brugada ECG pattern precipitated by acute pneumonia: A case report. *Cases J* 2009; 2: 73.
- Akhtar M, Goldschlager NF. Brugada electrocardiographic pattern due to tricyclic antidepressant overdose. *J Electrocardiol* 2006; 39: 336– 9.
- Al Aloul B, Adabag AS, Houghland MA, et al. Brugada pattern electrocardiogram associated with supratherapeutic phenytoin levels and the risk of sudden death. *Pacing Clin Electrophysiol* 2007; 30: 713– 715.
- Alhaj EK, Banning M, Vasaiwala SC, Santucci PA. Wolff-Parkinson-White, Brugada phenocopy, and flecainide toxicity: All in one patient. *Clin Case Rep*. 2019 Apr 23;7(5):1098-1102.
- Al-Sadawi M, Alkhawam H, Hartt A, Soleiman A, McFarlane SI. Cocaine-Induced Brugada Pattern. *Am J Med Case Rep*. 2019;7(4):59-61.
- Amin AS, Meregalli PG, Bardai A, et al. Fever increases the risk of cardiac arrest in the Brugada syndrome. *Ann Int Medicine* 2008; 149: 216– 218
- Amro M, Okoro KU, Mansoor K, Amro A, Okhumale P. Brugada Phenocopy Type 1 Secondary to Synthetic Cannabinoids. *Case Rep Cardiol*. 2020 Jun 23;2020:9425860.
- Ansari E, Cook JR. Profound hypothermia mimicking a Brugada type ECG. *J Electrocardiol* 2003; 36: 257– 260.
- Antzelevitch C, Brugada P, Borggrefe M, et al. Brugada syndrome: Report of the second consensus conference: endorsed by the Heart Rhythm Society and the European Heart Rhythm Association. *Circulation* 2005; 111: 659– 670.
- Antzelevitch C. The Brugada syndrome. Diagnostic criteria and cellular mechanisms. *Eur Heart J* 2001; 22: 356– 363.
- Arce M, Riera ARP, Femenia F, et al. Brugada electrocardiographic phenocopy in a patient with chronic Chagasic cardiomyopathy. *Cardiol J* 2010; 17: 525– 527.
- Babaliaros VC, Hurst JW. Tricyclic antidepressants and the Brugada syndrome: An example of Brugada waves appearing after the administration of desipramine. *Clin Cardiol* 2002; 25: 395– 398.
- Baranchuk A, Rosas F, Morillo CA. Enfermedad de Chagas en países desarrollados: mito o realidad (Chagas' disease in developed countries: myth or reality). In F Rosas, D Vanegas, M Cabrales (eds): *Enfermedad de Chagas*, Bogotá, Sociedad Colombiana de Cardiología. Bogota, Colombia, Sociedad Española de Cardiología, 2007, pp. 217– 219.G
- Baranchuk A, Simpson CS. Brugada syndrome coinciding with fever and pandemic (H1N1) influenza. *CMAJ* 2011; 183: 582.
- Baranchuk A, Antiperovitch P, Gottschalk BH. Brugada phenocopy due to hypokalemia. *J Family Med Prim Care*. 2018 Sep-Oct;7(5):1141-1142.

- Bebarta VS, Summers S. Brugada electrocardiographic pattern induced by cocaine toxicity. *Ann Emerg Med* 2007; 49: 827– 829.
- Bebarta VS, Waksman JC. Amitriptyline-induced Brugada pattern fails to respond to sodium bicarbonate. *Clin Toxicol* 2007; 45: 186– 188.
- Bernardo MH, Tiyyagura SR. A Case of Type I and II Brugada Phenocopy Unmasked in a Patient with Normal Baseline Electrocardiogram (ECG). *Am J Case Rep.* 2018 Jan 5;19:21-24.
- Bolognesi R, Tsialtas D, Vasini P, et al. Abnormal ventricular repolarization mimicking myocardial infarction after heterocyclic antidepressant overdose. *Am J Cardiol* 1997; 79: 242– 245.
- Bonnemeier H, Mauser W, Schunkert H. Images in cardiovascular medicine. Brugada-like ECG pattern in severe hypothermia. *Circulation* 2008; 118: 977– 978.
- Bramos D, Koutras K, Kollias G, et al. Cardiac amyloidosis and Brugada-like ECG pattern. *Int J Cardiol* 2009; 145: 249– 251.
- Brito MR, Miranda CES, Rabelo W, et al. Type 1 electrocardiographic Brugada pattern in a woman with Chagas disease: A case report. *Europace* 2010; 12: 1345– 1346.C
- Chiale PA, Przybylski J, Laiño RA, et al. Electrocardiographic changes evoked by ajmaline in chronic Chagas' disease without manifest myocarditis. *Am J Cardiol* 1982; 49: 14– 20.
- Chinushi M, Tagawa M, Izumi D, et al. Pilsicainide-induced ST segment depression in two patients with variant forms of Brugada-type electrocardiographic abnormalities. *Pacing Clin Electrophysiol* 2009; 32: 811– 815
- Chow BJ, Gollob M, Birnie D. Brugada syndrome precipitated by a tricyclic antidepressant. *Heart* 2005; 91: 651.
- Çinier G, Tse G, Baranchuk A. Brugada phenocopies: Current evidence, diagnostic algorithms and a perspective for the future. *Turk Kardiyol Dern Ars.* 2020 Mar;48(2):158-166.
- Cole JB, Sattiraju S, Bilden EF, et al. Isolated Tramadol overdose associated with Brugada ECG pattern. *Pacing Clin Electrophysiol* 2010 Oct 7 [E-pub ahead of print], doi: 10.1111/j.1540–8159.2010.02924.x.
- Copetti R, Proclemer A, Pillinini PP. Brugada-like ECG abnormalities during thioridazine overdose. *Br J Clin Pharmacol* 2005; 59: 608.
- Daccarett M, Freih M, Machado C. Acute cannabis intoxication mimicking Brugada-like ST segment abnormalities. *Int J Cardiol* 2007; 119: 235– 236.
- Daniel D Anselm, Jennifer M Evans, and Adrian Baranchuk, Brugada phenocopy: A new electrocardiogram phenomenon *World J Cardiol.* 2014 Mar 26; 6(3): 81–86. doi: 10.4330/wjc.v6.i3.81
- Dharanipradab M, Viswanathan S, Kumar GR, Krishnamurthy V, Stanley DD. Yellow phosphorus-induced Brugada phenocopy. *J Electrocardiol.* 2018 Jan-Feb;51(1):129-131.

- Doty B, Kim E, Phelps J, Akpunonu P. Pathophysiology of Hyperkalemia Presenting as Brugada Pattern on Electrocardiogram (ECG). *Am J Case Rep.* 2020 Jul 8;21:e923464.
- Dumaine R, Towbin JA, Brugada P, et al. Ionic mechanisms responsible for the electrocardiographic phenotype of the Brugada syndrome are temperature dependent. *Circ Res* 1999; 85: 803– 809.
- Eggebrecht H, Wieneke H, Erbel R. Image of the month. Brugada-like ECG pattern in a patient with isolated right ventricular infarction. *Herz* 2009; 34: 327.
- Elizari MV, Levi R, Acunzo RS, et al. Abnormal expression of cardiac neural crest cells in heart development: A different hypothesis for the etiopathogenesis of Brugada syndrome. *Heart Rhythm* 2007; 4: 359– 365.
- García-Niebla J, Bayés de Luna A, Baranchuk A. Multifactorial Brugada Phenocopy. *JAMA Intern Med.* 2018 Jun 1;178(6):872.
- Gavrielatos G, Letsas KP, Pappas LK, et al. Brugada electrocardiographic pattern induced during febrile state with marked leukocytosis. *Pacing Clin Electrophysiol* 2007; 30: 135– 136.
- Ghisoni E, Marandino L, Lombardi P, Bonzano A, Becco P, Aglietta M, Fizzotti M, Gay F, Rota Scalabrini D. Safe Use of Carfilzomib in a Patient with Multiple Myeloma and Intermittent Type 1 Brugada ECG Pattern: A Case Report. *Acta Haematol.* 2019 Sep 25:1-5.
- Goldgran-Toledano D, Sideris G, Kevorkian JP. Overdose of cyclic antidepressants and the Brugada syndrome. *N Engl J Med* 2002; 346: 1591– 1592.
- Guamán Valdivieso C, Ferrando-Castagnetto F, Marichal P, Acquistapace F, Trujillo P, Baranchuk A. Brugada phenocopy secondary to extensive inferior wall ischemia induced by coronary steal effect. *Ann Noninvasive Electrocardiol.* 2020 Jan;25(1):e12689.
- Habuchi Y, Furukawa T, Tanaka H, et al. Ethanol inhibition of Ca²⁺ and Na⁺ currents in the guinea-pig heart. *Eur J Pharmacol* 1995; 292: 143– 149. 36
- Heckle M, Alsafwah S, Agarwal M. Multifactorial Brugada Phenocopy-Reply. *JAMA Intern Med.* 2018 Jun 1;178(6):873.
- Hisamatsu K, Kusano KF, Morita H, et al. Relationships between depolarization abnormality and repolarization abnormality in patients with Brugada Syndrome: Using body surface signal-averaged electrocardiography and body surface maps. *J Cardiovasc Electrophysiol* 2004; 15: 870– 876.
- Irani F, Kasmani R, Kanjwal Y. Hyperkalemia and cocaine induced dynamic Brugada-type electrocardiogram. *Eur J Emerg Med* 2009; 17: 113– 115,
- Itoh E, Suzuki K, Tanabe Y. A case of vasospastic angina presenting Brugada-type ECG abnormalities. *Jpn Circ J* 1999; 63: 493– 495.

- Janse MJ, Kleber AG. Electrophysiological changes and ventricular arrhythmias in the early phase of regional myocardial ischemia. *Circ Res* 1981; 49: 1069– 1081.
- Jesús Monterrubio-Villar 1, David Llinares-Moya 1Brugada Phenocopy Induced by a Lethal Methanol Intoxication *Eur J Case Rep Intern Med*. 2020 Jan 1;7(2):001374. doi: 10.12890/2020_001374.
- Junttila MJ, Gonzalez M, Lizotte E, et al. Induced Brugada-type electrocardiogram, a sign for imminent malignant arrhythmias. *Circulation* 2008; 117: 1890– 1893.
- Kaiser E, Pastore CA, Moffa PJ. Ebstein's anomaly with Brugada-like electrocardiogram pattern: a critical view. *Cardiology* 2010; 17: 415.
- Kalra S, Iskandar SB, Duggal S, et al. Fever-induced ST-segment elevation with a Brugada syndrome type electrocardiogram. *Ann Intern Med* 2008; 148: 82– 84.
- Karakitso D, Patrianakos A, Poularas J, et al. Brugada-like electrocardiographic pattern unmasked by fever in a critical care patient. *Acta Anaesthesiol Scand* 2006; 50: 1038– 1039.
- Kataoka H. Electrocardiographic patterns of the Brugada Syndrome in 2 young patients with pectus excavatum. *J Electrocardiol* 2002; 35: 169– 171.
- Keller DI, Rougier JS, Kucera JP, et al. Brugada syndrome and fever: Genetic and molecular characterization of patients carrying SCN5A mutations. *Cardiovasc Res* 2005; 67: 510– 519.
- Khalil Y, Siddique SM, Hoang NT. A Brugada-pattern electrocardiogram and adrenal insufficiency. *Ann Intern Med* 2010; 153: 849– 851.
- Kim YH, Lim HE, Kim SH, et al. Brugada-like ST-segment abnormalities associated with myocardial involvement of hematologic diseases. *Pacing Clin Electrophysiol* 2008; 31: 761– 764.
- Kiran HS, Ravikumar YS, Jayasheelan MR, et al. Brugada like pattern in ECG with drug overdose. *J Assoc Physicians India* 2010; 58: 120– 122.
- Kovacic JC, Kuchar DL. Brugada pattern electrocardiographic changes associated with profound electrolyte disturbance. *Pacing Clin Electrophysiol* 2004; 27: 1020– 1023.
- Kurisu S, Inoue I, Kawagoe T. Brugada-like electrocardiographic pattern due to hyperkalemia. *Clin Cardiol* 2009; 32: E23.
- Kusaka K, Yamakawa J, Kawaura K, et al. Brugada-like electrocardiographic changes during influenza infection. *J Int Med Res* 2003; 31: 244– 246.
- Kutsuzawa D, Arimoto T, Watanabe T, et al. Persistent abnormal value of late potential in Brugada syndrome associated with hypokalemia. *Ann Noninvasive Electrocardiol* 2011; 16:104– 106.

- Lancini D, Shetty R. Tension pneumothorax: a Brugada phenocopy and ST-elevation myocardial infarction mimic. *Eur Heart J*. 2020 Jun 14;41(23):2163.
- Lazaros G, Lazarou E, Tousoulis D. Brugada phenocopy in a patient undergoing pericardiocentesis for a large idiopathic pericardial effusion. *J Electrocardiol*. 2019 Jan 7:S0022-0736(18)30853-7.
- Levine M, Lovecchio F. Diphenhydramine-induced Brugada pattern. *Resuscitation* 2010; 81: 503– 504.
- Lippolis A, Buzzi MP, Romano IJ, Facchini C, Gentile F. Syncope in a patient with acute pulmonary embolism and Brugada Type-2 ECG pattern: Brugada phenocopy or Brugada syndrome? *J Electrocardiol*. 2019 May-Jun;54:90-92.
- Littmann L, Monroie MH, Taylor III L, et al. The hyperkalemic Brugada sign. *J Electrocardiol* 2007; 40: 53– 59.
- Littmann L, Monroe MH, Svenson RH. Brugada-type electrocardiographic pattern induced by cocaine. *Mayo Clin Proc* 2000; 75: 845– 849.
- Li Y, Liu T, Tse G, Tao L. Brugada phenocopy following coronary artery bypass graft surgery. *J Electrocardiol*. 2020 Mar-Apr;59:134-139.
- Lopez-Barbeito B, Llis M, Delgado V, et al. Diphenhydramine overdose and Brugada sign. *Pacing Clin Electrophysiol* 2005; 28: 730– 732.
- Makaryus JN, Verbsky J, Schwartz S, et al. Fever associated with gastrointestinal shigellosis unmasks probable Brugada syndrome. *Case Report Med* 2009;492031.
- Meert A, Vermeersch N, Beckers R, et al. Brugada-like ECG pattern induced by tricyclic antidepressants. *Eur J Emerg Med* 2010; 17: 325– 327.
- Mehta S, Parameswaran AC, Greenspan A, et al. Hypercalcemia due to rhabdomyolysis mimicking Brugada syndrome. *Pacing Clin Electrophysiol* 2009; 32: e14– 15.
- Meregalli P, Ruijter J, Hofman N, et al. Diagnostic value of flecainide testing in unmasking SCN5A-related Brugada syndrome. *J Cardiovasc Electrophysiol* 2006; 17: 857– 864.
- Mok N, Tong C, Yuen H. Concomitant-acquired long QT and Brugada syndromes associated with indapamide-induced hypokalemia and hyponatremia. *Pacing Clin Electrophysiol* 2008; 31: 772– 775.
- Monterrubio-Villar J, Llinares-Moya D. Brugada Phenocopy Induced by a Lethal Methanol Intoxication. *Eur J Case Rep Intern Med*. 2020 Jan 1;7(2):001374.
- Nademanee K, Veerakul G, Chandanamattha P, et al. Prevention of ventricular fibrillation episodes in Brugada syndrome by catheter ablation over the anterior right ventricular outflow tract epicardium. *Circulation* 2011; 123: 1270– 1279.
- Nagase S, Kusano KF, Morita H, et al. Epicardial electrogram of the right ventricular outflow tract in patients with the Brugada syndrome. *J Am Coll Cardiol* 2002; 39: 1992– 1995.

- Nakazato Y, Kurata T, Yamaguchi H. ST segment elevation in the precordial leads mimicking Brugada syndrome. *Heart* 2000; 83: 216.
- Nakazato Y, Ohmura T, Shimada I, et al. Brugada-like precordial ST elevation on ECG by anterior mediastinal infective mass lesion. *Indian Pacing Electrophysiol J* 2003; 3: 184.
- Nayyar S, Nair M. Brugada pattern in toxic myocarditis due to severe aluminum phosphide poisoning. *Pacing Clin Electrophysiol* 2009; 32: e16– 17.
- Nishida K, Fujiki A, Mizumaki K, et al. Canine model of Brugada syndrome using regional epicardial cooling of the right ventricular outflow tract. *J Cardiovasc Electrophysiol* 2004; 15: 936– 941
- Noda T, Shimizu W, Taguchi A, et al. ST-segment elevation and ventricular fibrillation without coronary spasm by intracoronary injection of acetylcholine and/or ergonovine maleate in patients with Brugada syndrome. *J Am Coll Cardiol* 2002; 40: 1841– 1847.
- Ortega-Carnicer J, Benezet J, Calderon-Jimenez P, et al. Hypothermia-induced Brugada-like electrocardiogram pattern. *J Electrocardiol* 2008; 41: 690– 692.
- Ortega-Carnicer J, Benezet J, Ruiz-Lorenzo F, et al. Transient Brugada-type electrocardiographic abnormalities in renal failure reversed by dialysis. *Resuscitation* 2002; 55: 215– 219.
- Ortega-Carnicer J, Bertos-Polo J, Gutierrez-Tirado C. Aborted sudden death, transient Brugada pattern, and wide QRS dysrhythmias after massive cocaine ingestion. *J Electrocardiol* 2001; 34: 345– 349.
- Ott P, Freund NS. Brugada-pattern EKG in a febrile patient. *J Emerg Med* 2007; 33: 281– 282.
- Ozben B, Caymaz O, Erdogan O. Fever-induced precordial ST-segment elevation in a young man. *Turk Kardiyoloji Dernegi Arsivi* 2010; 38: 35– 37.
- Ozeke O, Aras D, Geyik B, et al. Brugada-type electrocardiographic pattern induced by fever. *Indian Pacing Electrophysiol J* 2005; 5: 146– 148.
- Ozeke O, Selcuk MT, Topaloglu S, et al. Brugada-like early repolarisation pattern associated with acute pericarditis. *EMJ* 2006; 23: e64
- Palaniswamy C, Selvaraj DR, Chugh T, et al. Brugada electrocardiographic pattern induced by amitriptyline overdose. *Am J Ther* 2010; 17: 529– 532.
- Patanè S, Marte F. Revelation of Brugada electrocardiographic pattern during a febrile state associated with acute myocardial infarction. *Int J Cardiol* 2010; 144: e1– 4.

- Paul G, Yusuf S, Sharma S. Unmasking of the Brugada syndrome phenotype during the acute phase of amiodarone infusion. *Circulation* 2006; 114: e489– 491.
- Phillips N, Priestley M, Denniss AR, et al. Brugada-type electrocardiographic pattern induced by epidural bupivacaine. *Anesth Analg* 2003; 97: 264– 267.
- Pirotte MJ, Mueller JG, Poprawski T. A case report of Brugada-type electrocardiographic changes in a patient taking lithium. *Am J Emerg Med* 2008; 26: e1– 3.
- Porres JM, Brugada J, Urbistondo V, et al. Fever unmasking the Brugada syndrome. *Pacing Clin Electrophysiol* 2002; 25: 1646– 1648.
- Postema PG, Wolpert C, Amin AS, et al. Drugs and Brugada syndrome patients: Review of the literature, recommendations and an up-to-date website (<http://www.brugadadrugs.org>). *Heart Rhythm* 2009; 6: 1335– 1341.
- R Gunaseelan 1, M Sasikumar 2, K Aswin 2, Shirshendu Dhar 2, N Balamurugan 2, Vivekanandan Pillai 2 Brugada phenocopy induced by consumption of yellow oleander seeds - A case report. *J Electrocardiol.* 2020 Jul 24;62:107-109. doi: 10.1016/j.jelectrocard.2020.07.014. Online ahead of print. *J Electrocardiol.* 2020 Jul 24;62:107-109. doi: 10.1016/j.jelectrocard.2020.07.014. Online ahead of print.
- Ramsaroop K, Seecheran R, Seecheran V, Persad S, Giddings S, Mohammed B, Seecheran NA. Suspected hyponatremia-induced Brugada phenocopy. *Int Med Case Rep J.* 2019 Feb 27;12:61-65.
- Rangaraj R, Moorthy N, Patil SS, et al. Brugada-type electrocardiographic pattern induced by electrocution. *Indian Pacing Electrophysiol J* 2009; 9: 56– 59.
- Refaat MM, Hotait M, London B. Genetics of sudden cardiac death. *Curr Cardiol Rep.* 2015;17:606.
- Refaat MM, Hotait M, Scheinman MM. Brugada syndrome. *Card Electrophysiol Clin.* 2016;8:239-245.
- Riera ARP, Uchida AH, Schapachnik E, et al. Propofol infusion syndrome and Brugada syndrome electrocardiographic phenocopy. *Cardiol J* 2010; 17: 130– 135.
- Riezzo I, Centini F, Neri M, et al. Brugada-like EKG pattern and myocardial effects in a chronic propofol abuser. *Clin Toxicol* 2009; 47: 358– 363. Cr54
- Robertson KE, Martin TN, Rae AP. Brugada-pattern ECG and cardiac arrest in cocaine toxicity: Reading between the white lines. *Heart* 2010; 96: 643– 644.
- Roberts-Thomson KC, Teo KS, Young GD. Drug-induced Brugada syndrome with ST-T wave alternans and long QT. *Intern Med J* 2007; 37: 199– 200.

- Rollin A, Maury P, Guilbeau-Frugier C, et al. Transient ST elevation after ketamine intoxication: A new cause of acquired Brugada ECG pattern. *J Card Elect* 2010; 22: 91– 94.
- Rouleau F, Asfar P, Boulet S, et al. Transient ST segment elevation in right precordial leads induced by psychotropic drugs: Relationship to the Brugada syndrome. *J Cardiovasc Electrophysiol* 2001; 12: 61– 65.
- Rudnik-Schoneborn S, Schaupp M, Lindner A, et al. Brugada-like cardiac disease in myotonic dystrophy type 2: Report of two unrelated patients. *Eur J Neurol* 2010; 18: 191– 194.
- Sasaki A, Nakazato Y. Brugada-like electrocardiogram detected after reconstructive operation for oesophageal cancer. *Europace* 2010; 12: 1542
- Saura D, Garcia-Alberola A, Carrillo P, et al. Brugada-like electrocardiographic pattern induced by fever. *Pacing Clin Electrophysiol* 2002; 25: 856– 859.
- Sawhney V, Thomas G, Webster P, et al. Resolution of Brugada-pattern ECG after withdrawal of the selective serotonin reuptake inhibitor paroxetine. *Heart* 2010; 96: 1165– 1166.
- Sharma AN, Hexdall AH, Chang EK, et al. Diphenhydramine-induced wide complex dysrhythmia responds to treatment with sodium bicarbonate. *Am J Emerg Med* 2003; 21: 212– 215.
- Sheikh M, Kanjwal K, Kasmani R, et al. Simultaneous ST-segment elevation in inferior and precordial leads following ingestion of a lethal dose of desipramine: A novel Brugada-like EKG pattern. *J Interv Card Electrophysiol* 2010; 28: 35– 38.
- Shim A, Seecheran R, Seecheran V, Persad S, Sreenivasan S, Henry R, Seecheran NA. Suspected Brugada Phenocopy Secondary to Coronary Slow Flow. *Case Rep Cardiol.* 2019 Dec 6;2019:9027029.
- Shimada M, Miyazaki T, Miyoshi S, et al. Sustained monomorphic ventricular tachycardia in a patient with Brugada syndrome. *Jpn Circ J* 1996; 60: 364– 370.
- Shimizu W. Acquired forms of the Brugada syndrome. *J Electrocardiol* 2005; 38: 22– 25.
- Siniorakis E, Arvanitakis S, Psatheris G, et al. Recurrence of fever-related Brugada pattern and right ventricular functional correlates. *Int J Cardiol* 2010; 148: 370– 372.
- Sreenivasan S, Monaghan M, Baranchuk A. Multifactorial Brugada Phenocopy. *JAMA Intern Med.* 2018 Jun 1;178(6):872-873.
- Stirnimann G, Severine P, Abriel H, et al. Brugada syndrome ECG provoked by the selective serotonin reuptake inhibitor fluvoxamine. *Europace* 2010; 12: 282– 283.
- Suzuki T, Kohsaka S. Brugada-type electrocardiographic changes in a febrile patient of African descent. *Am J Med Sci* 2006; 332: 97– 99.

- Tada H, Sticherling C, Oral H, Morady F. Brugada syndrome mimicked by tricyclic antidepressants overdose. *J Cardiovasc Electrophysiol* 2001; 12: 275.
- Takagi M, Aihara N, Kuribayashi S, et al. Localized right ventricular morphological abnormalities detected by electron-beam computed tomography represent arrhythmogenic substrates in patients with the Brugada syndrome. *Eur Heart J* 2001; 22: 1032– 1041.C
- Tamene A, Sattiraju S, Wang K, et al. Brugada-like electrocardiography pattern induced by severe hyponatraemia. *Europace* 2010; 12: 905– 907.
- Tanawuttiwat T, Harindhanavudhi T, Bhan A, et al. Hyperkalemia-induced Brugada pattern: An unusual manifestation. *J Cardiovasc Med* 2010; 11: 285– 287.
- Tarin N, Farre J, Rubio JM, et al. Brugada-like electrocardiographic pattern in a patient with a mediastinal tumor. *Pacing Clin Electrophysiol* 1999; 22: 1264– 1266.
- Teijeiro R, Garro HA, Acunzo RS, et al. Recording of high V1-V3 precordial leads may be essential to the diagnosis of Brugada syndrome during the ajmaline test. *J Cardiovasc Pharmacol Ther* 2006; 11: 153– 155.
- Tomcsanyi J, Simor T, Papp L. Haemopericardium and Brugada-like ECG pattern in rheumatoid arthritis. *Heart* 2002; 87: 234.
- Tomcsanyi J, Zsoldos A, Bozsik B. Brugada-like acute myocardial infarction. *Heart* 2003; 89: 1199.
- Tsai CF, Wu DJ, Lin MC, et al. A Brugada-pattern electrocardiogram and thyrotoxic periodic paralysis. *Ann Intern Med* 2010; 153: 848– 849.
- Tsarouhas K, Papalexis P, Kafantaris I, et al. Electrocardiographic findings compatible with Brugada syndrome in a patient with febrile respiratory infection. *Hippokratia* 2010; 14: 221– 223.
- Unlu M, Bengi F, Amasyali B, et al. Brugada-like electrocardiographic changes induced by fever. *EMJ* 2007; 24: e4.Cr
- Vernooij K, Delhaas T, Cremer Ol, et al. Electrocardiographic changes predicting sudden death in propofol-related infusion syndrome. *Heart Rhythm* 2006; 3: 131– 137.
- Wahbi K, Fressart V, Bécane HM, et al. High prevalence of Brugada syndrome in patients with Steinert's disease. *Medgen* 2009; 21: 423.
- Weiner JB, Haddad EV, Raj SR. Recovery following propofol-associated Brugada electrocardiogram. *Pacing Clin Electrophysiol* 2010; 33: e39– 42r
- Wilde AA, Postema PG, DiDiego JM, et al. The pathophysiological mechanism underlying Brugada syndrome depolarization versus repolarization. *J Mol Cell Cardiol* 2010; 49: 543– 553.
- Wright D, Salehian O. Brugada-type electrocardiographic changes induced by long-term lithium use. *Circulation* 2010; 122: e418– 419.

- Xu G, Gottschalk BH, Anselm DD, Benditt DG, Maheshwari A, Sreenivasan S, Shama RA, Dendramis G, Barajas-Martínez H, Rubio Campal JM, Aznaurov SG, Baranchuk A. Relation of the Brugada Phenocopy to Hyperkalemia (from the International Registry on Brugada Phenocopy). *Am J Cardiol*. 2018 Mar 15;121(6):715-717.
- Xu G, Gottschalk BH, Pérez-Riera A, Barbosa-Barros R, Dendramis G, Carrizo AG, Agrawal S, Bayés de Luna A, Jastrzębski M, Tomcsányi J, Baranchuk A. Link between Brugada phenocopy and myocardial ischemia: Results from the International Registry on Brugada Phenocopy. *Pacing Clin Electrophysiol*. 2019 Jun;42(6):658-662.
- Yan GX, Antzelevitch C. Cellular basis for the Brugada syndrome and other mechanisms of arrhythmogenesis associated with ST-Segment elevation. *Circulation* 1999; 100: 1660– 1666.
- Yu M, Zhang Q, Huang X, Zhao X. Type 1 Brugada phenocopy in a patient with acute pericarditis. *J Electrocardiol*. 2018 Nov-Dec;51(6):1121-1123.
- Zakka P, Refaat MM. Brugada pattern, Brugada phenocopy, or Brugada syndrome: That is the question. *Pacing Clin Electrophysiol*. 2019 Jul;42(7):777-778.
- Zampi G, Pergolini A, Pontillo D, Sommariva L. A tricky electrocardiogram: cocaine induced Brugada type 2 phenocopy. *Am J Cardiovasc Dis*. 2020 Jun 15;10(2):131-135.
- Zhuo M, Li J, Lecker SH. Brugada phenocopy induced by hyperkalemia. *Kidney Int*. 2019 Feb;95(2):471.