

Apnea del Sueño:

...el corazón sufre hasta cuando dormimos...



Adrián Baranchuk
Associate Professor of Medicine and Physiology
Queen's University
Kingston, Ontario, Canada

**Simposio Hospital Cosme Argerich
Octubre 2010**

Apnea del Sueño y Enfermedad Cardiovascular

1. Definiciones y manifestaciones clínicas
2. Epidemiología
3. Fisiopatología
4. El *link* entre AS y el aparato cardiovascular
5. Métodos de estudio
6. Breve resumen del tratamiento
7. Líneas de investigación

Apnea del Sueño

Definiciones

1. **AS**: presencia de apneas e hipopneas durante el sueño, sintomáticas por despertares abruptos (“arousals”), ronquidos y somnolencia diurna
2. **Apnea**: interrupción del flujo aéreo, con reducción de la saturación de $O_2 \geq 4\%$
 - ▶ **Central**: deterioro de los centros nerviosos que regulan la respiración
 - ▶ **Obstruktiva**: reducción del flujo aéreo en la vía respiratoria alta
3. **Hipopnea**: disminución $\geq 30\%$ del flujo aéreo o excursión tóraco-abdominal,

con reducción de la saturación de $O_2 \geq 4\%$
4. **AHI (Apnea/hipopnea index)**: cantidad de apneas e hipopneas por hora

Apnea del Sueño: Fisiopatología



Progress in
Cardiovascular
Diseases

Vol. 41, No. 5
March/April 1999

Pathogenesis of Obstructive Sleep Apnea

M. Safwan Badr

Mecanismos de obstrucción de la vía aérea durante el sueño:

1. Presión intraluminal subatmosférica

2. Angostamiento espiratorio

3. Reducción ventilatoria motora

4. Teoría mixta: central + obstructiva

El link entre AS y morbilidad cardiovascular

Cardiovascular Morbidity in Obstructive Sleep Apnea

J. Woodrow Weiss, Sandrine H. Launois, Amit Anand, and Erik Garpestad

Progress in Cardiovascular Diseases, Vol. 41, No. 5 (March/April), 1999: pp 367-376

1. Cambios fisiológicos durante el sueño

- a. En fase NREM: caída de TA y FC (10-20%): ↓ del VM y RPT, ↓ Tono Simpático en fase 4 de NREM
- b. En fase REM: vasoconstricción + fluctuación del VM y FC

2. Cambios agudos durante AS


- a. FC: existen 2 patrones de comportamiento
 - Bradicardia al comienzo de la apnea, aceleración durante la AS (quimiorreceptores) y pico de aceleración al final de la apnea y al despertar (*más frecuente*)
 - Bradicardia progresiva a lo largo de la apnea
- b. TA: Incremento al final de la apnea (> desaturación, ↑ tono simpático por despertar abrupto)
- c. Volumen minuto: ↓ volumen de eyección durante la apnea acompañado de presión negativa intrapleurales

Mecanismos fisiopatológicos de morbilidad cardiovascular

Síndrome de apneas-hipopneas durante el sueño y corazón

Joaquín Terán Santos^a, M. Luz Alonso Álvarez^a, José Cordero Guevara^b, José María Ayuela Azcárate^c y José María Monserrat Canal^d

Rev Esp Cardiol. 2006;59(7):718-24



Resumiendo...

Cardiovascular Morbidity in Obstructive Sleep Apnea

J. Woodrow Weiss, Sandrine H. Launois, Amit Anand, and Erik Garpestad

Progress in Cardiovascular Diseases, Vol. 41, No. 5 (March/April), 1999: pp 367-376

AS



Fluctuaciones de TA, FC y Volumen minuto

Desaturación O₂



Morbilidad cardiovascular

Cambios agudos

- Arritmias
- Infarto de miocardio
- ACV

Cambios crónicos

- ICC
- HTA
- HTP
- HVI

Apnea del Sueño: AS & Arritmias

1. Bradiarritmias

- Paro sinusal
- Bloqueo AV

2. Taquiarritmias

- Fibrilación auricular
- Arritmias ventriculares

Qué se ha demostrado de todo esto?


Apnea del Sueño: AS & Arritmias

Association of Nocturnal Arrhythmias with Sleep-disordered Breathing

The Sleep Heart Health Study

AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE VOL 173 2006

	SDB, % (n = 228)	No SDB, % (n = 338)	Pearson's χ^2 p Value
Ventricular Arrhythmias			
Premature Ventricular Contraction ($\geq 5/h$)	35.1	21.3	0.0003
Bigeminy	14.0	8.0	0.02
Trigeminy	9.2	5.6	0.10
Quadrigeminy	11.8	5.9	0.01
Nonsustained ventricular tachycardia	5.3	1.2	0.004
Complex ventricular ectopy*	25	14.5	0.002
Supraventricular Arrhythmias			
Premature atrial contraction ($\geq 5/h$) [†]	33.8	24.3	0.001
Atrial fibrillation	4.8	0.9	0.003
Supraventricular tachycardia	14.9	14.5	0.89
Conduction Delay Arrhythmias			
Sinus pause (≥ 3 s)	11.0	8.6	0.34
First-degree atrioventricular block	25.0	22.5	0.49
Second-degree atrioventricular block type 1	1.8	0.3	0.07
Second-degree atrioventricular block type 2	2.2	0.9	0.20
Intraventricular conduction delay	8.9	5.3	0.11




La asociación entre AS y FA es altamente significativa. AS también se asocia con Arritmias ventriculares.

Sleep Apnea: SA & AF


Association of Atrial Fibrillation and Obstructive Sleep Apnea

Apoor S. Gami, MD; Gregg Pressman, MD; Sean M. Caples, MD; Ravi Kanagala, MD;
Joseph J. Gard, BS; Diane E. Davison, RN, MA; Joseph F. Malouf, MD; Naser M. Ammash, MD;
Paul A. Friedman, MD; Virend K. Somers, MD, PhD



	OR	95% CI	OR and 95% CI	P
Body mass index	1.11	1.06-1.16		<0.0001
Neck circ	1.02	0.97-1.07		0.439
Hypertension	1.27	1.01-1.61		0.039
Diabetes mellitus	1.23	0.96-1.57		0.104
Atrial fibrillation	2.19	1.40-3.42		0.0006

It's time to wake up!: sleep apnea and cardiac arrhythmias




Apnea del Sueño: AS & Arritmias

It's time to wake up!: sleep apnea and cardiac arrhythmias

- (i) Impaired autonomic nervous control has been demonstrated in patients with OSA, manifesting as increased sympathetic tone and/or decreased parasympathetic tone. Decreased baroreflex sensitivity, reduced vagal input, and impairment of the parasympathetic components of the heart rate variability have been demonstrated in patients with OSA.^{22,23}
- (ii) A persistent increase in sympathetic tone, as occurring in OSA, has been shown to generate abnormal electrical remodelling of the atrium, facilitating supraventricular arrhythmias, and AF in particular.²⁴ Specifically, electrical remodelling may create some degree of interatrial block, contributing to the genesis of atrial arrhythmias.²⁵
- (iii) A strong association between OSA and hypertension has been reported extensively.^{6,26} As well, the association between hypertension and AF is well recognized.^{27,28} Although purely speculative, the link

Understanding the association between sleep apnea & cardiac arrhythmias



Adrian Branchuk MD FACC¹, Christopher S. Simpson MD FACC FRCPC¹, Damian P. Redfearn MB ChB MRCP¹,
Kevin Michael MD¹, Mike Fitzpatrick MD FRCPI, FRCPC, D.ABSM²



Apnea del Sueño: AS & Arritmias

Bradycardia during Sleep Apnea


CHARACTERISTICS AND MECHANISM



- Bradycardia durante AS
- Cuanto mayor es la apnea mayor es la bradicardia
- No bradicardia en ausencia de AS

Truths and Lies from the Polysomnography ECG Recording: An Electrophysiologist Perspective

El registro ECG del polisomnógrafo es una sola derivación y a veces puede confundirnos. En este caso el paciente fue enviado por un bloqueo 2:1 y en verdad se trata de extrasístoles del tracto de salida del VD.



Apnea del Sueño: Tratamiento con Marcapasos

Physiologic Pacing in Patients With Obstructive Sleep Apnea

A Prospective, Randomized Crossover Trial

Andrew D. Krahn, MD,* Raymond Yee, MD,* Mark K. Erickson, BSc,† Toby Markowitz, BSEE,†
Lorne J. Gula, MD,* George J. Klein, MD,* Allan C. Skanes, MD,* Charles F. P. George, MD,*
Kathleen A. Ferguson, MD*

Parameter	Control	AAI Pacing	Difference (95% CI)	p Value
Apnea hypopnea index	42.1 ± 20.7	38.6 ± 20.5	-3.4 (-9.3 to 2.5)	0.23
Heart rate (beats/min)	65 ± 7	77 ± 1	12 (8 to 16)	0.001
Time in bed (min)	437 ± 45	417 ± 108	-19 (-87 to 48)	0.55
Minimum SaO ₂ (%)	77 ± 11	75 ± 10	-2 (-9 to 4)	0.38
% time SaO ₂ <90%	3.5 ± 4.3	3.8 ± 6.0	0.3 (-1.4 to 2.1)	0.70
Apnea hypopnea duration (s)	28.4 ± 4.7	33.1 ± 11.2	4.7 (0.7 to 8.7)	0.11
Circulatory time (s)	25.5 ± 4.4	23.4 ± 3.2	-2.1 (-4.7 to 0.4)	0.09

El marcapaseo secuencial no demostró beneficios
en pacientes con AS

Atrial overdrive pacing: 12 RCT desde 2002...

Effects of physiological cardiac pacing on sleep-disordered breathing in patients with chronic bradydysrhythmias *Psychiatry and Clinical Neurosciences* (2001), 55, 257–258

Atrial Overdrive Pacing for the Obstructive Sleep Apnea–Hypopnea Syndrome

N Engl J Med 2005;353:2568–77.

Physiologic Pacing in Patients

- Los beneficios no son claros...

Atrial overdrive pacing compared to CPAP in patients with obstructive sleep apnoea syndrome

European Heart Journal (2005) 26, 2568–2575

Overdrive atrial pacing does not improve obstructive sleep apnoea syndrome

Eur Respir J 2005; 25: 343–347


Effect of atrial overdrive pacing on obstructive sleep apnea in patients with systolic heart failure ☆

Sleep Medicine 8 (2007) 31–36

Atrial overdrive pacing in sleep apnoea: a meta-analysis

Adrian Baranchuk^{1*}, Jeff S. Healey², Christopher S. Simpson¹, Damian P. Redfearn¹, Carlos A. Morillo², Stuart J. Connolly², and Michael Fitzpatrick¹


Review: OSA_AOP_Baranchuk
Comparison: 01 Apnea-Hypopnea Index
Outcome: 01 AHI



Reducción del AHI en 5 puntos:
Significativo estadísticamente pero...
Sin relevancia clínica!!!

Sleep Disordered Breathing And Ventricular Arrhythmias: Mechanisms and Implications

	SA (N=26)	NO SA (N=106)	p
Incidence of Therapies			
Appropriate (%)	31	17	0.09
Inappropriate (%)	3.8	9.4	NS
Time to First Appropriate Therapy (months $\pm$ SD)	8.0 $\pm$ 5.2	11.89 $\pm$ 5.9	0.12
Incidence of Arrhythmias			
NSVT (%)	34.6	50.0	NS
SVT (%)	26.9	30.1	NS



Apnea del Sueño: Enfermedad coronaria

Increased incidence of coronary artery disease in sleep apnoea: a long-term follow-up

Eur Respir J 2006;28:596-602


(n=308)

Seguimiento: 7 años

AS: AHI ≥ 30

BASAL	OSA	Non-OSA	p-value*
Subjects n	105	203	
Age yrs	51.8 $\pm$ 8.9	47.6 $\pm$ 10.2	<0.001
BMI kg·m ⁻²	28.6 $\pm$ 4.0	25.9 $\pm$ 3.7	<0.001
SBP mmHg	138.4 $\pm$ 16.3	130.1 $\pm$ 17.9	<0.001
DBP mmHg	82.2 $\pm$ 9.4	78.6 $\pm$ 10.3	0.003
OD events·h ⁻¹	94.1 $\pm$ 86.0	8.9 $\pm$ 8.3	<0.001
ODI events·h ⁻¹	17.8 $\pm$ 15.1	1.6 $\pm$ 1.7	<0.001
Sa _o ₂ ,min %	80.5 $\pm$ 7.8	88.6 $\pm$ 4.2	<0.001
Males	91 (86.7)	154 (75.9)	0.026
Hypertension	30 (28.6)	33 (16.3)	0.011
Diabetes mellitus	2 (1.9)	2 (1.0)	NS
Smokers	35 (34.0)	80 (41.5)	NS
CAD incidence	17 (16.2)	11 (5.4)	0.003
Cardiovascular death	8 (7.6)	1 (0.5)	<0.001

	β	RR (95% CI)	p-value
Constant	-22.84		
OSA at baseline	1.53	4.60 (1.83–11.6)	0.001
Time since baseline yrs	0.19	1.21 (1.01–1.45)	0.043
SBP at baseline mmHg	0.04	1.03 (1.01–1.05)	<0.001
Current age yrs	0.06	1.06 (1.02–1.11)	0.007
Sa _o ₂ ,min at baseline %	0.11	1.11 (1.02–1.22)	0.019



Apnea del Sueño: Enfermedad coronaria

Sleep-disordered Breathing and Cardiovascular Disease

Cross-sectional Results of the Sleep Heart Health Study

AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE VOL 163 2001

(n=6424)

AHI dividido en cuartiles

Análisis multivariado y ajustado para covariables

	Quartile				p Value†
	I	II	III	IV	
Coronary heart disease					
Full model	1.0	1.01 (0.77–1.32)	1.20 (0.92–1.57)	1.22 (0.93–1.59)	<u>0.08</u>
Parsimonious model	1.0	0.92 (0.71–1.20)	1.20 (0.93–1.54)	1.27 (0.99–1.62)	<u>0.004</u>
Heart failure					
Full model	1.0	1.19 (0.56–2.53)	1.96 (0.99–3.90)	2.20 (1.11–4.37)	0.008
Parsimonious model	1.0	1.13 (0.54–2.39)	1.95 (0.99–3.83)	2.38 (1.22–4.62)	0.002
Stroke					
Full model	1.0	1.24 (0.76–2.01)	1.38 (0.86–2.83)	1.55 (0.96–2.50)	0.06
Parsimonious model	1.0	1.15 (0.72–1.83)	1.42 (0.91–2.21)	1.58 (1.02–2.46)	0.03


Mecanismos relacionados

1. Hipertensión Arterial
2. Hiperactividad simpática diurna
3. Hipoxemia
4. ↑ agregabilidad plaquetaria
5. Ruptura aguda de placa
6. Hipertensión Pulmonar


Apnea del Sueño: Enfermedad coronaria

Benefits of obstructive sleep apnoea treatment in coronary artery disease: a long-term follow-up study[☆]

- (n=54)
- Pacientes on AS + Enf. Coronaria
- CPAP/ cirugía vs. no tratamiento
- End-point compuesto: mortalidad cardiovascular, evento isquémico, hospitalización por ICC, revascularización



Tratamiento	No tratamiento
6/25 (24%)	17/29 (58%)
HR 0.24 (95% CI 0.09 – 0.62)	
P<0.01	





A quien le preocupa este tema...?

Sleep Apnea and Cardiovascular Disease

An American Heart Association/American College of Cardiology
Foundation Scientific Statement From the American Heart Association
Council for High Blood Pressure Research Professional Education
Committee, Council on Clinical Cardiology, Stroke Council, and Council on
Cardiovascular Nursing Council

*In Collaboration With the National Heart, Lung, and Blood Institute National Center on Sleep
Disorders Research (National Institutes of Health)*





AGOSTO 2008 (On-LINE)


Apnea del Sueño: Accidente Cerebrovascular

Investigating the Relationship Between Stroke and Obstructive Sleep Apnea

Debieramos reformular el
CHADS2 score?




Dyken ME. Stroke 1996;27:401-407



AS & Insuficiencia cardíaca

Mecanismo de interacción

Rev Esp Cardiol. 2006;59(7):718-24



AS & Insuficiencia cardíaca

Escenarios clínicos

1. Disfunción ventricular izquierda

(n=47)

Fey < 40%

AHI > 15

Resultados

1. AS: 55% de los ptes

2. Más frecuente en CAD

Central Sleep Apnea in Left Ventricular Dysfunction Prevalence and Implications for Arrhythmic Risk

Paola A. Lanfranchi, MD; Virend K. Somers, MD, PhD; Alberto Braghiroli, MD; Ugo Corra, MD; Ermanno Eleuteri, MD; Pantaleo Giannuzzi, MD

Background—The prevalence and characteristics of sleep-disordered breathing in patients with asymptomatic left ventricular (LV) dysfunction are unknown. Therefore, we evaluated the prevalence of sleep-disordered breathing in patients with LV dysfunction without overt heart failure and tested the hypothesis that sleep-disordered breathing is linked to greater hemodynamic and autonomic impairment.

Methods and Results—We studied 47 patients with LV ejection fractions $\leq 40\%$ without any history of heart failure. Central sleep apnea (CSA), as defined by an apnea-hypopnea index $\geq 15/h$, was present in 26 patients (55%), 17 (36%) of whom had severe CSA (apnea-hypopnea index $\geq 30/h$). Obstructive sleep apnea was evident in 5 patients (11%). The prevalence and severity of CSA were higher in patients with ischemic cardiomyopathy than in patients with nonischemic cardiomyopathy ($P < 0.05$). Exercise tolerance and echocardiographic indices of systolic and diastolic function were similar in patients without CSA, with mild CSA, and with severe CSA. Heart rate variability was markedly depressed in patients with CSA ($P < 0.05$). Patients with severe CSA also had a higher incidence of nonsustained ventricular tachycardia ($P = 0.05$).


Conclusions—CSA is highly prevalent in patients with asymptomatic LV dysfunction. The severity of CSA may not be related to the severity of hemodynamic impairment. Severe CSA is associated with impaired cardiac autonomic control and with increased cardiac arrhythmias. (*Circulation*. 2003;107:727-732.)

Key Words: sleep ■ nervous system, autonomic ■ tachyarrhythmias ■ heart failure

In patients with overt heart failure, there is a high prevalence of nocturnal periodic breathing with central apneas (central sleep apnea: CSA).¹⁻⁴ CSA is associated with increased arrhythmic risk⁵ and may indicate increased mortality in heart failure.⁴ Autonomic responses to CSA may contribute to the adverse prognosis in these patients.^{2,5} Patients with left ventricular (LV) dysfunction without heart failure also have neurohumoral activation and are at risk for progression to

Methods

We prospectively studied consecutive patients referred to the Cardiology Department of the Medical Center of Rehabilitation, Veruno, Italy, between January 1999 and December 2000 who were found to have LV systolic dysfunction due to either ischemic or nonischemic cardiomyopathy. Patients were referred for one of the following: (1) functional evaluation of asymptomatic LV dysfunction, (2) evaluation of chest pain, or (3) rehabilitation after myocardial infarction or cardiac surgery. They were eligible



AS & Insuficiencia cardíaca

Escenarios clínicos

2. Insuficiencia cardíaca congestiva

(n=450)

Fey $27.3 \pm 15\%$

NYHA II: 62%, NYHA III: 34%

FA: 15%

AHI > 20

Resultados

1. AS (AHI>20): 53%

2. Más frec. en FA ($p < 0.01$)

Am J Respir Crit Care Med 1999;160:1101-06

Risk Factors for Central and Obstructive Sleep Apnea in 450 Men And Women with Congestive Heart Failure

DON D. SIN, FABIA FITZGERALD, JOHN D. PARKER, GARY NEWTON, JOHN S. FLORAS, and T. DOUGLAS BRADLEY

Sleep Research Laboratory of the Toronto Rehabilitation Institute and the Departments of Medicine, The Toronto Hospital and Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada

In previous analyses of the occurrence of central (CSA) and obstructive sleep apnea (OSA) in patients with congestive heart failure (CHF), only men were studied and risk factors for these disorders were not well characterized. We therefore analyzed risk factors for CSA and OSA in 450 consecutive patients with CHF (382 male, 68 female) referred to our sleep laboratory. Risk factors for CSA were male gender (odds ratio [OR] 3.50; 95% confidence interval [CI], 1.39 to 8.84), atrial fibrillation (OR 4.13; 95% CI 1.53 to 11.14), age > 60 yr (OR 2.37; 95% CI 1.35 to 4.15), and hypocapnia ($P_{CO_2} < 38$ mm Hg during wakefulness) (OR 4.33; 95% CI 2.50 to 7.52). Risk factors for OSA differed by gender: in men, only body mass index (BMI) was significantly associated with OSA (OR for a BMI > 35 kg/m², 6.10; 95% CI 2.86 to 13.00); whereas, in women, age was the only important risk factor (OR for age > 60 yr, 6.04; 95% CI 1.75 to 20.0). We conclude that historical information, supplemented by a few simple laboratory tests may enable physicians to risk stratify CHF patients for the presence of CSA or OSA, and the need for diagnostic polysomnography for such patients. Sin DD, Fitzgerald F, Parker JD, Newton G, Floras JS, Bradley TD. Risk factors for central and obstructive sleep apnea in 450 men and women with congestive heart failure. *AM J RESPIR CRIT CARE MED* 1999;160:1101-1106.

Obstructive sleep apnea (OSA) is an important risk factor for the development of hypertension, angina pectoris, myocardial infarction, and cor pulmonale (1-4). More recent data suggest that sleep apnea can also lead to the progression of cardiac dysfunction in patients with chronic congestive heart failure (CHF) (5, 6). This adverse effect on cardiac function probably arises from repetitive apneas causing arterial oxygen desaturation, excessive stimulation of the sympathetic nervous system, and increases in systemic blood pressure (5-7).

The presence of central sleep apnea (CSA) in patients with CHF is associated with a significantly increased risk for death and cardiac transplantation (8, 9). In addition, fragmentation of sleep architecture by frequent arousals can also lead to the development of excessive daytime sleepiness and fatigue in CHF patients with either OSA or CSA (5, 6, 10). Treatments specifically aimed at OSA and CSA in patients with CHF have been shown to improve cardiovascular function and clinical status. For example, continuous positive airway pressure (CPAP) has been shown to alleviate both OSA and CSA, and to improve left ventricular ejection fraction (LVEF), decrease urinary and plasma norepinephrine concentrations, and improve

symptoms of heart failure (5, 11, 12). Oxygen also alleviates CSA and reduces nocturnal urinary norepinephrine concentrations (13).

A recent report suggests that both CSA and OSA are common in the CHF population (14). The clinical characteristics of patients with CSA appear to differ from those with OSA, probably reflecting important differences in the underlying pathophysiologies of these two breathing disorders. In a study of men with CHF, Javaheri and coworkers found that those with OSA were heavier and had a higher prevalence of snoring than those with CSA or no sleep apnea (14). Patients with CSA, on the other hand, had a lower LVEF. However, because only 81 patients (of whom only nine had OSA) were evaluated in their study, other factors that distinguish risk for OSA from those for CSA or no sleep-related breathing disorder (SBD) may have escaped detection. More importantly, because only men were studied, risk factors for sleep apnea in women with CHF remain unknown. Indeed, risk factors for CSA in women either with or without CHF have not been reported. The purpose of our study, therefore, was twofold: first, to determine the overall risk factors for CSA and OSA in a large group of 450 patients with CHF referred to our sleep laboratory; and second, to determine whether there are differences in risk factors for CSA and OSA between men and women with chronic, stable CHF.

METHODS

Subjects

We conducted a retrospective analysis of 450 consecutive patients with CHF, referred to the Toronto Rehabilitation Institute Sleep Research Laboratory between July 1987 and November 1998. All patients were referred to the sleep laboratory by cardiologists. The crite-

(Received in original form March 1, 1999 and in revised form May 19, 1999)

Supported by an operating grant from the Ontario Thoracic Society.

D. Sin is supported by a research fellowship from the Alberta Heritage Foundation for Medical Research. G. Newton is supported by a Research Scholarship from the Heart and Stroke Foundation of Ontario, and J. Floras is supported by a Career Investigator award from the Heart and Stroke Foundation of Ontario.

Correspondence and requests for reprints should be addressed to T. Douglas Bradley, M.D., E5 12-421 The Toronto Hospital (General Division), 200 Elizabeth St., Toronto, ON, M5G 2C4 Canada. E-mail: douglas.bradley@utoronto.ca

Am J Respir Crit Care Med Vol 160, pp 1101-1106, 1999
Internet address: www.atsjournals.org

AS & Insuficiencia cardíaca

Escenarios clínicos

3. Disfunción diastólica

(n=20)
NYHA II-III
AHI > 10

Resultados

1. AS (AHI>10): 55%
2. Tiempo de desaceleración
($p<0.05$)

Chest 1997;111:1488-93



clinical investigations

Prevalence of Sleep-Disordered Breathing in Diastolic Heart Failure*


Joseph Chan, MBBS; John Sanderson, MD; Wilson Chan, MBBS;
Christopher Lai, DM, FCCP; Dominic Choy, MBBS; Alice Ho, MBBS; and
Roland Leung, MD, FCCP

Objective: Sleep-disordered breathing (SDB) is common in congestive heart failure. While isolated diastolic heart failure (DHF) accounts for up to a third of all cases of congestive heart failure, the prevalence of SDB in DHF is unknown. We aim to determine the prevalence and characteristics of SDB in a group of patients with symptomatic DHF.

Methods: Twenty subjects with symptomatic DHF (New York Heart Association class II or III) and isolated diastolic dysfunction on echocardiography were assessed with lung function tests, modified sleep and health questionnaire, and overnight polysomnography. Significant SDB was defined as an apnea/hypopnea index (AHI) >10.

Results: Thirteen female and seven male subjects (mean age, 65 ± 6.0 years; mean body mass index (BMI), 28 ± 3.2) were evaluated, of whom 17 (85%) had a diagnosis of hypertension. Overall sleep quality was poor, with fragmentation and frequent arousals associated with respiratory events. Fifty-five percent of the patients had significant SDB, mainly obstructive apneas. BMI and the prevalence of hypertension were similar in patients with and without SDB. The deceleration time, an index of diastolic dysfunction, was more prolonged in the group with SDB (236 ± 40 ms vs 282 ± 31 ms; $p<0.05$). As a group, a lower minimum percentage arterial oxygen saturation during sleep, but not the AHI was associated with more severe degree of diastolic dysfunction on echocardiogram, including a lower ratio between the early peak transmitral flow velocity and the late peak atrial systolic velocity ($\rho=0.57$; $p<0.05$) and a prolonged isovolumic relaxation time ($\rho=-0.54$; $p<0.05$).

Conclusions: SDB is common in patients with DHF. Patients with DHF and SDB may be associated with worse diastolic dysfunction than those without SDB, although a causal relationship remains to be established. (Chest 1997;111:1488-93)



Sleep-disordered breathing (SDB) is common in congestive heart failure (CHF). The prevalence of SDB in CHF is 45% in

*From the Department of Medicine, Prince of Wales Hospital, Shatin, NT, Hong Kong. Manuscript received 1997.

Reprint requests: Dr. Joseph Chan, Prince of Wales Hospital, Shatin, NT, Hong Kong.

AS & Terapia de Resincronización Cardíaca

Cardiac Resynchronization Therapy Improves Central Sleep Apnea and Cheyne-Stokes Respiration in Patients With Chronic Heart Failure

Anil-Martin Sinha, MD, DPHIL,* Erik C. Skobel, MD,* Ole-Alexander Breithardt, MD,* Christine Norra, MD,† Kai U. Markus, MD,* Christian Breuer, MD,* Peter Hanrath, MD, FESC, FACC,* Christoph Stellbrink, MD, FESC*

Aachen, Germany

(n=24)

Non-randomized study

Ambulatory polygraph

		All Patients (n = 24)	CSA (n = 14)	No CSA (n = 10)	p Value (CSA vs. No CSA)
Cardiorespiratory polygraph					
Duration of recording (h)	Pre	8 ± 2	8 ± 2	8 ± 3	NS
	CRT	8 ± 3	8 ± 3	8 ± 2	NS
AHI (per h)	Pre	11.9 ± 11.7 (1-42)	19.2 ± 10.3 (9-42)	1.7 ± 0.7 (1-3)	0.00001
	CRT	3.3 ± 3.8* (0-12)	4.6 ± 4.4* (0-12)	1.5 ± 1.6 (0-4)	NS
SaO ₂ min (%)	Pre	88 ± 5 (70-92)	84 ± 5 (70-90)	90 ± 2 (88-92)	<0.05
	CRT	90 ± 2* (85-93)	89 ± 2* (85-90)	91 ± 1 (90-93)	NS
PSQI	Pre	7.5 ± 4.2 (2-14)	10.4 ± 1.6 (8-14)	2.4 ± 0.5 (2-4)	<0.05
	CRT	3.5 ± 2.2* (2-10)	3.9 ± 2.4* (2-10)	2.6 ± 0.9 (2-4)	NS

1. Significant reduction of AHI (pre CRT vs post CRT)
2. Significant increment of SaO₂
3. Significant reduction of PSQI (Pittsburgh Sleep Quality Index)

AS & Terapia de Resincronización Cardíaca

Improvement in Cheyne-Stokes respiration following cardiac resynchronisation therapy


J.Y. Gabor, D.A. Newman, V. Barnard-Roberts, V. Korley, I. Mangat, P. Dorian and P.J. Hanly

1. uncontrolled
2. observational
3. small sample

TABLE 3 Sleep apnoea following 6 months of cardiac resynchronisation therapy (CRT)

	Pre-CRT	Post-CRT
Subjects n	10	10
Total AHI [†] ·h ⁻¹	42.7 ± 9.1	30.8 ± 18.7
CSR events ·h ⁻¹ *	30.6 ± 14.0	15.3 ± 16.5
Obstructive AHI ·h ⁻¹	9.7 ± 12.3	11.8 ± 10.2
Mean Sa _{o2} ,awake %	93.6 ± 1.0	93.9 ± 0.7
Mean Sa _{o2} ,sleep %	93.3 ± 1.1	93.3 ± 1.1
Mean PCO _{2,tc} ,awake mmHg	42.1 ± 4.1	43.6 ± 4.9
Mean PCO _{2,tc} ,sleep mmHg	42.2 ± 3.9	44.9 ± 4.4
PB cycle length s	63.6 ± 16.2	59.1 ± 9.6
LFCT s	43.0 ± 12.4	35.6 ± 6.8 [#]

Data are presented as mean ± SD, unless otherwise stated. AHI: apnoea-hypopnoea index; CSR: Cheyne-Stokes respiration; Sa_{o2}: arterial oxygen saturation; PCO_{2,tc}: transcutaneous partial pressure of carbon dioxide; PB: periodic breathing; LFCT: lung-to-finger circulation time. [†]: includes mixed apnoeas in addition to CSR and obstructive apnoeas. *: p < 0.05 versus pre-CRT; [#]: p = 0.09.



6/10 have significantly reduced AHI
Two thirds have reduced CSR

Apnea del Sueño: AS & Hipertensión Arterial

PROSPECTIVE STUDY OF THE ASSOCIATION BETWEEN SLEEP-DISORDERED BREATHING AND HYPERTENSION

PAUL E. PEPPARD, PH.D., TERRY YOUNG, PH.D., MARI PALTA, PH.D., AND JAMES SKATRUD, M.D.

NEJM 2000;342:1378-84

TABLE 3. ADJUSTED ODDS RATIOS FOR HYPERTENSION AT A FOLLOW-UP SLEEP STUDY, ACCORDING TO THE APNEA-HYPOPNEA INDEX AT BASE LINE.*

BASE-LINE APNEA–HYPOPNEA INDEX	HTA		HTA + BMI	tabaco
	ODDS RATIO, ADJUSTED FOR BASE-LINE HYPER- TENSION STATUS	ODDS RATIO, ADJUSTED FOR BASE-LINE HYPER- TENSION STATUS AND NONMODIFIABLE RISK FACTORS (AGE AND SEX)	ODDS RATIO, ADJUSTED FOR BASE-LINE HYPER- TENSION STATUS, NON- MODIFIABLE RISK FAC- TORS, AND HABITUS (BMI AND WAIST AND NECK CIRCUMFERENCE)	ODDS RATIO, ADJUSTED FOR BASE-LINE HYPER- TENSION STATUS, NON- MODIFIABLE RISK FAC- TORS, HABITUS, AND WEEKLY ALCOHOL AND CIGARETTE USE
	odds ratio (95% confidence interval)			
0 events/hr†	1.0	1.0	1.0	1.0
0.1–4.9 events/hr	1.66 (1.35–2.03)	1.65 (1.33–2.04)	1.42 (1.14–1.78)	1.42 (1.13–1.78)
5.0–14.9 events/hr	2.74 (1.82–4.12)	2.71 (1.78–4.14)	2.03 (1.29–3.19)	2.03 (1.29–3.17)
≥15.0 events/hr	4.54 (2.46–8.36)	4.47 (2.37–8.43)	2.89 (1.47–5.69)	2.89 (1.46–5.64)
P for trend‡	<0.001	<0.001	<0.002	<0.002

(n=709)

AS & Hipertensión Arterial

Grandes estudios poblacionales

Wisconsin Sleep Study¹: (n=709), seg. hasta 8 años

Sleep Heart Health Study²: (n=6841), seg. 3 años

Wisconsin Sleep Study			Sleep Heart Health Study		
Punto de corte IAH	Prevalencia HTA (%)	OR* (IC del 95%)	Punto de corte IAH	Prevalencia HTA (%)	OR* (IC del 95%)
0	17	1	< 1,5	43	1
0,1-4,9	28	1,39 (1,04-1,84)	1,5-4,9	53	1,07 (0,91-1,26)
5-14,9	48	1,92 (1,09-3,39)	5-14,9	59	1,20 (1,01-1,42)
≥ 15	60	2,66 (1,13-6,25)	15-29,9	62	1,25 (1,00-1,56)
			≥ 30	67	1,37 (1,03-1,83)

A mayor severidad de AS, la asociación con HTA se incrementa



La pregunta es: ¿**existe causalidad?**

1. N Engl J Med 2000;342:1378-84
2. JAMA 2000;283:1829-36

AS & Hipertensión Arterial Fisiopatología

Fisiopatología de la hipertensión asociada al síndrome de apnea obstructiva del sueño: Evidencia de estudios clínicos y modelos animales de hipoxia crónica intermitente

Rev Med Chile 2007; 135: 1333-1342



Rey et al. 2008

AS & Hipertensión Pulmonar

La controversia continúa...

Pulmonary Artery Hypertension and Sleep-Disordered Breathing*

ACCP Evidence-Based Clinical Practice Guidelines
Chest 2004;126:72-77

Source	AHI or RDI, h	BMI (or %IBW)†	%COPD	Definition of PH	PAP Measurement Method	Prevalence of PH, No./Total (%)	Factors With Significant Association with PH
Alchanatis et al. ¹⁰ 2001	> 15	NR	NR	mPAP > 20 mm Hg	RH cath	6/29 (20.7)	Age (p < 0.05) BMI (p < 0.02) Daytime PO ₂ (p < 0.05) Daytime PCO ₂ (p < 0.001) FVC and FEV ₁ (p < 0.001) Body weight (p = 0.0002) BMI (p = 0.002) FEV ₁ (p < 0.001)†
Apprill et al. ¹¹ 1991	NR	NR	NR	mPAP ≥ 20 mm Hg	RH cath	9/46 (20)	FEV ₁ /FVC (p < 0.001) PO ₂ (p < 0.001) AHI (p < 0.001) Mean nocturnal SaO ₂ (p < 0.001)† BMI (p < 0.01) FEV ₁ /FVC < 65% PCO ₂ †
Bady et al. ¹² 2000	> 5	NR	Excluded	mPAP > 20 mm Hg	RH cath	12/44 (27)	PO ₂ (p < 0.001) PCO ₂ (p < 0.001) TLC, FEV ₁ , FEV ₁ /FVC, %FVC (p < 0.001) AHI, apnea index (p < 0.001) Daytime SaO ₂ (p < 0.001) PO ₂ (p < 0.001)† PCO ₂ (p < 0.001) FEV ₁ (p < 0.001)† FEV ₁ /FVC (p = 0.03) SaO ₂ (p = 0.01)
Chaouat et al. ¹³ 1996	> 20	NR	NR	mPAP ≥ 20 mm Hg	RH cath	37/200 (17)	BMI (p < 0.01) Awake pH (p < 0.05) PaCO ₂ (p < 0.05) VC (p < 0.01) No difference in FEV ₁ /FVC, FRC %pred
NS					RH cath	13/65 (20) 31/65 (48) with exercise	NR
NS					Doppler echo	11/22 (50)	No difference in age, BMI, smoking history
NS					RH cath	18/92 (20)	PCWP (p < 0.005) Sleep time with SpO ₂ < 90% (p < 0.05) FVC, FVC %pred, FEV ₁ , FEV ₁ /FVC (p < 0.001) PaO ₂ (p < 0.001) PaCO ₂ (p < 0.001)
NS					Doppler echo	8/37 (22)	Minimum SpO ₂ (p = 0.051)

RECOMENDACIONES

1. En la evaluación de ptes. con HTP, evaluar AS es mandatorio (evidencia:baja, rédito:pobre, clase C)
2. En la evaluación de ptes. con HTP por AS, la Polisomnografía está indicada (evidencia:opinion de experto, rédito:intermedio, clase B)
3. En el manejo de ptes. con AS evaluación de HTP de rutina NO es recomendada (evidencia:baja, rédito:no, clase I)
4. En ptes con AS e HTP, CPAP reduce la presión pulmonar (pero NO la normaliza) (evidencia:baja, rédito:pobre, clase C)

pulmonary artery pressures, while 2 studies relied on estimates derived echocardiographically. In general, PH associated with OSA was mild. The average mPAP

in each of these studies was < 30 mm Hg; in most, it was < 25 mm Hg. Prevalence estimates of PH ranged from 17 to 53%.

Apnea del Sueño & Hipertrofia ventricular izquierda

Prevalence of Left Ventricular Hypertrophy in Persons With and Without Obstructive Sleep Apnea

(Cardiology in Review 2006;14: 170–172)

- (n=53)
- RDI > 5, severo > 30
- HVI: ♀ 110 g/m² / ♂ 134 g/m²

Variable	OSA (n = 40)	No OSA (n = 13)	P
Age (years)	51 ± 8	50 ± 8	NS
Men	24 (60%)	3 (23%)	<0.025
Women	16 (40%)	10 (77%)	<0.025
Hypertension	17 (43%)	5 (38%)	NS
Diabetes mellitus	9 (23%)	3 (23%)	NS
Body mass index ≥30 kg/m ²	22 (55%)	7 (54%)	NS
Coronary artery disease	10 (25%)	4 (31%)	NS

	LVH
Moderate or severe OSA (n = 27)	21 (78%)
Mild OSA (n = 13)	6 (46%)
No OSA (n = 13)	3 (23%)

P < 0.001 comparing moderate or severe OSA with no OSA.
P < 0.05 comparing moderate or severe OSA with mild OSA.

El link entre AS e HVI podría ser la *hipertensión arterial*, sin embargo, esta presente en chicos y adolescentes con AS

Apnea del Sueño & Sistema Nervioso Autónomo

• (n=60)

• 3

• A

• H

• -

• -

• A

HRV Data	Mild OSAS (n=19)	Severe OSAS (n=17)	Control (n=24)	Cardiac Autonomic Activity			Apnea
				P1	P2	P3	
SDNN (ms)	124.94	120.38	131.05	<0.05	<0.01	NS	obstructive vided them 9) had mild dings were or age, sex, Holter moni- variability. In ular arrhyth- patients with vere OSAS, e similar. In ut in severe AS patients e lower. The up II than in
SDANN (ms)	112.29	108.63	126.95	<0.05	<0.05	NS	
RMSSD (ms)	35.11	31.54	44.16	NS	<0.05	NS	
Triangular index*	33.88	32.33	36.90	NS	NS	NS	
Total power (ms ²)	1,405.35	1,695.63	1,290.42	<0.01	<0.01	NS	patients with absence of the severity
Ultra low frequency (ms ²)	165.24	137.54	119.05	<0.01	<0.05	NS	
Very low frequency (ms ²)	582.77	633.71	510.21	<0.05	<0.05	NS	
Low frequency (ms ²)	432.12	476.58	411.16	<0.05	<0.01	<0.05	
High frequency (ms ²)	341.12	332.50	351.47	<0.05	<0.01	NS	ring sleep. ¹ ypnoea, is sleep ² and sleep apnea arterial hy- so an asso-
Low-frequency/high-frequency ratio	1:4	1:4	1:3	<0.05	<0.01	<0.05	

- ↓ tono parasimpático
- ↑ tono simpático

Address for reprints:
Mustafa Aydin, MD,
Zonguldak Karaelmas
Universitesi Tıp Fakültesi,
Kardiyoloji Anabilim Dalı,
67600 Kozlu, Zonguldak,
Turkey

E-mail:
drmustafaaydin@hotmail.com

© 2004 by the Texas Heart*
Institute, Houston

measurements has also been considered useful in the screening of sleep apnea patients.¹² However, the relationship between HRV and the cardiac autonomic dysfunction of OSAS is not completely understood. Therefore, we prospectively investigated by time-dependent and spectral analysis, HRV in patients with OSAS and in a control group.

Patients and Methods



Study Population

In our study, the existence of OSAS was determined by clinical findings and by the apnea hypopnea index (AHI): ≥ 5 episodes of apnea or hypopnea per hour of sleep. We enrolled 36 patients with OSAS and divided them according to the apnea hypopnea index (AHI) into 2 groups: Group I (n=19) had mild OSAS (AHI <20)

(Tex Heart Inst J 2004;31:132-6)

Apnea del Sueño & Regulación de la VFC

Determinants of heart rate variability in obstructive sleep apnea syndrome during wakefulness and sleep



3 mecanismos

1. Estímulo vagal de receptores pulmonares
2. Control medular de los centros respiratorios
3. Sensibilidad barorrefleja


↓ **significativa aferencia vagal**
($P < 0.04$)

↓ **significativa SBR**
($P < 0.03$)

Apnea del Sueño & Regulación de la VFC

Dynamic time-varying analysis of heart rate and blood pressure variability in cats exposed to short-term chronic intermittent hypoxia

Am J Physiol Regul Integr Comp Physiol 295: R28–R37, 2008.



Hipoxia Cronica Intermitente (8 Hr-4 días):

1. ↓ Sensibilidad Barorrefleja
2. ↓ Total VFC (LF predominante)

Apnea del Sueño: Diagnóstico: Polisomnografía


Diagnostic Techniques in Obstructive Sleep Apnea

Charles Frederick Petersen George

Progress in Cardiovascular Diseases, Vol. 41, No. 5 (March/April), 1999: pp 355-366



Diferencias entre Apnea Obstructiva y Apnea Central





Apnea del Sueño: Tratamiento con CPAP

Continuous Positive Airway Pressure Therapy for Treating Sleepiness in a Diverse Population With Obstructive Sleep Apnea

Results of a Meta-analysis *Arch Intern Med.* 2003;163:565-571

- 12 estudios randomizados
- End point: reducción en la escala de Epworth



> beneficio
en ptes con
AHI > 30

CPAP mejor que placebo (OR 2.94, $p < 0.001$)
Heterogeneidad – $p < 0.001$

Algunas Conclusiones

Si desea contactarme: adribaran@hotmail.com

1. La Apnea del Sueño es altamente prevalente.
2. Se asocia con enfermedades cardiovasculares como ICC, HTA, Obesidad y Síndrome Metabólico, Stroke, Enfermedad Coronaria, Disfunción Autonómica y Arritmias.
3. Conocer los mecanismos fisiopatológicos permite encarar tratamientos multifactoriales (reducir peso, controlar HTA, prevenir arritmias e infartos).
4. Su reconocimiento permite tratarla con C-PAP
5. La evidencia indica que el marcapaseo auricular rápido NO debe ser universalmente recomendado.
6. En pacientes con Apnea de origen CENTRAL, el tratamiento de resincronización cardíaca mejora notablemente los índices de apnea. Debe ser parte de la consideración al recomendar esta terapia